




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 403/14, 413/14, 417/14	A1	(11) International Publication Number: WO 97/47618 (43) International Publication Date: 18 December 1997 (18.12.97)												
<p>(21) International Application Number: PCT/US97/09888</p> <p>(22) International Filing Date: 6 June 1997 (06.06.97)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>60/019,487</td> <td>10 June 1996 (10.06.96)</td> <td>US</td> </tr> <tr> <td>9614190.8</td> <td>5 July 1996 (05.07.96)</td> <td>GB</td> </tr> <tr> <td>60/023,312</td> <td>31 July 1996 (31.07.96)</td> <td>US</td> </tr> <tr> <td>9617897.5</td> <td>28 August 1996 (28.08.96)</td> <td>GB</td> </tr> </table> <p>(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): LIVERTON, Nigel, J. [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). BUTCHER, John, W. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). CLAREMON, David, A. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). BILODEAU, Mark, T. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p>		60/019,487	10 June 1996 (10.06.96)	US	9614190.8	5 July 1996 (05.07.96)	GB	60/023,312	31 July 1996 (31.07.96)	US	9617897.5	28 August 1996 (28.08.96)	GB	<p>(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
60/019,487	10 June 1996 (10.06.96)	US												
9614190.8	5 July 1996 (05.07.96)	GB												
60/023,312	31 July 1996 (31.07.96)	US												
9617897.5	28 August 1996 (28.08.96)	GB												

(54) Title: SUBSTITUTED IMIDAZOLES HAVING CYTOKINE INHIBITORY ACTIVITY



(57) Abstract

Compounds represented by formula (I) are disclosed. AR represents an aromatic group containing 6-10 atoms; and (a) represents a 4 to 10 membered non-aromatic heterocycle containing at least one N atom, and optionally containing 1-2 additional N atoms and 0-1 O or S atom. A pharmaceutical composition is also included. Methods of treating cancer and cytokine mediated diseases are also included.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

-1-

TITLE OF THE INVENTION**SUBSTITUTED IMIDAZOLES HAVING CYTOKINE INHIBITORY ACTIVITY****5 BACKGROUND OF THE INVENTION**

 The present invention relates to substituted imidazole compounds which have cytokine inhibitory activity. Cytokine mediated diseases and cytokine inhibition, suppression and antagonism are used in the context of diseases or conditions in which excessive or unregulated production or activity of one or more cytokines occurs. Examples of
10 cytokines which are effected typically include Interleukin-1 (IL-1), Interleukin-6 (IL-6), Interleukin-8 (IL-8) and Tumor Necrosis Factor (TNF).

 Interleukin-1 (IL-1) and Tumor Necrosis Factor (TNF) are
15 produced by a variety of cells which are involved in immunoregulation and other physiological conditions.

 There are many disease states in which IL-1 is implicated. Examples are rheumatoid arthritis, osteoarthritis, endotoxemia, toxic shock syndrome, acute and chronic inflammatory diseases, such as the
20 inflammatory reaction induced by endotoxin or inflammatory bowel disease; tuberculosis, atherosclerosis, muscle degeneration, cachexia, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis and acute synovitis. Recent evidence also links IL-1 activity to diabetes.

 Interleukin-1 has been demonstrated to mediate a variety of biological activities thought to be important in immunoregulation and other physiological conditions. [See, e.g., Dinarello et al., Rev. Infect. Disease, 6, 51 (1984)]. The known biological activities of IL-1 include the activation of T helper cells, induction of fever, stimulation of
25 prostaglandin or collagenase production, neutrophil chemotaxis, induction of acute phase proteins and the suppression of plasma iron levels.
30

 Excessive or unregulated tumor necrosis factor (TNF) production or activity has been implicated in mediating or exacerbating

-2-

5 rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis, and other arthritic conditions, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, adult respiratory distress syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcosis, bone resorption diseases, reperfusion injury, graft v. host rejection, allograft rejections, fever and myalgia due to infection, cachexia secondary to infection or malignancy, cachexia secondary to acquired immune deficiency syndrome (AIDS), AIDS related complex (ARC), keloid formation, scar tissue formation, 10 Crohn's disease, ulcerative colitis and pyresis.

Monokines, such as TNF, have also been shown to activate HIV replication in monocytes and/or macrophages [See Poli, *et al.*, Proc. Natl. Acad. Sci., 87:782-784 (1990)], therefore, inhibition of monokine production or activity aids in limiting HIV progression. TNF has been 15 implicated in various roles with other viral infections, such as the cytomegalovirus (CMV), influenza virus and the herpes virus.

Interleukin-6 (IL-6) is a cytokine effecting the immune system and hematopoiesis. It is produced by several mammalian cell types in response to agents such as IL-1, and is correlated with disease states such 20 as angiofollicular lymphoid hyperplasia.

Interleukin-8 (IL-8) is a chemotactic factor first identified and characterized in 1987. Many different names have been applied to IL-8, such as neutrophil attractant/activation protein-1 (NAP-1), monocyte derived neutrophil chemotactic factor (MDNCF), neutrophil 25 activating factor (NAF), and T-cell lymphocyte chemotactic factor. Like IL-1, IL-8 is produced by several cell types, including mononuclear cells, fibroblasts, endothelial cells and ketainocytes. Its production is induced by IL-1, TNF and by lipopolysaccharide (LPS). IL-8 stimulates a number of cellular functions *in vitro*. It is a chemoattractant for neutrophils, T- 30 lymphocytes and basophils. It induces histamine release from basophils. It causes lysozomal enzyme release and respiratory burst from neutrophils, and it has been shown to increase the surface expression of Mac-1 (CD11b/CD 18) on neutrophils without *de novo* protein synthesis.

-3-


There remains a need for compounds which are useful in treating cytokine mediated diseases, and as such, inhibit, suppress or antagonize the production or activity of cytokines such as IL-1, IL-6, IL-8 and TNF.

5

SUMMARY OF THE INVENTION

The present invention addresses a compound represented by formula I:

10




I

or a pharmaceutically acceptable salt thereof, wherein:

15

X and X' each independently represent $-(CH_2)_m-Y-(CH_2)_n-$, wherein m and n represent integers within the range of from 0 - 4, such that the sum of m and n is from 0 - 6; Y represents a member selected from the group consisting of: a direct bond; O; $S(O)_y$, with y equal to 0, 1 or 2; $NR^{q'}$, with $R^{q'}$ as defined below; $C(O)$; $OC(O)$; $C(O)O$; $SO_xNR^{q'}$ with x equal to 1 or 2 and $R^{q'}$ as defined below; $NR^{q'}SO_x$; $C(O)NR^{q'}$ and $NR^{q'}C(O)$;

20



represents a 4 to 10 membered non-aromatic heterocycle containing at least one N atom, and optionally containing 1-2 additional N atoms and 0-1 O or S atom;

-4-

R^x represents H, C₁₋₆ alkyl(R^q)₃, C₃₋₈ cycloalkyl, OC₁₋₆ alkyl(R^q)₃ or C(O)C₁₋₆ alkyl(R^q)₃;


each R independently represents a member selected from the group consisting of: halo; hydroxy; C₁₋₆ alkyl(R^q)₃; OC₁₋₆ alkyl(R^q)₃; C₃₋₈ cycloalkyl(R^q)₃; CN; CONH₂; CONHC₁₋₆ alkyl(R^q)₃; CON(C₁₋₆ alkyl(R^q)₃)₂; NH₂; NHC₁₋₆ alkyl(R^q)₃; NHC₃₋₈ cycloalkyl; N(C₁₋₆ alkyl(R^q)₃)₂; CON(C₃₋₈ cycloalkyl)(C₁₋₆ alkyl(R^q)₃); CO₂H; CO₂C₁₋₆ alkyl(R^q)₃; C(O)C₁₋₆ alkyl(R^q)₃; aryl(R^q)₃; heterocyclyl(R^q)₃; heteroaryl(R^q)₃; CF₃; SH; NO₂; NHSO₂C₁₋₆alkyl(R^q)₃, NHSO₂aryl(R^q)₃, NHSO₂heteroaryl(R^q)₃; N(R^{q'})C(O)C₁₋₆ alkyl(R^q)₃; NR^{q'}C(O)NH(C₁₋₆ alkyl(R^q)₃); C₂₋₄ alkenyl(R^q)₂₋₃ and C₂₋₄ alkynyl(R^q)₁₋₃;

each R'' independently represents a member selected from the group consisting of: halo; hydroxy; C₁₋₆ alkyl(R^q)₃; OC₁₋₆ alkyl(R^q)₃; C₃₋₈ cycloalkyl(R^q)₃; CN; CONH₂; CONHC₁₋₆ alkyl(R^q)₃; CON(C₁₋₆ alkyl(R^q)₃)₂; NH₂; NHC₁₋₆ alkyl(R^q)₃; NHC₃₋₈ cycloalkyl; N(C₁₋₆ alkyl(R^q)₃)₂; CON(C₃₋₈ cycloalkyl)(C₁₋₆ alkyl(R^q)₃); CO₂H; CO₂C₁₋₆ alkyl(R^q)₃; C(O)C₁₋₆ alkyl(R^q)₃; aryl(R^q)₃; heterocyclyl(R^q)₃; heteroaryl(R^q)₃; CF₃; SH; NO₂; SO_yC₁₋₆ alkyl(R^q)₃, with y as defined above; SO₂NH₂; SO₂NHC₁₋₆ alkyl(R^q)₃; SO₂N(C₁₋₆ alkyl(R^q)₃)₂; NHSO₂C₁₋₆alkyl(R^q)₃, NHSO₂aryl(R^q)₃, NHSO₂heteroaryl(R^q)₃; N(R^{q'})C(O)C₁₋₆ alkyl(R^q)₃; NR^{q'}C(O)NH(C₁₋₆ alkyl(R^q)₃); C₂₋₄ alkenyl(R^q)₂₋₃ and C₂₋₄ alkynyl(R^q)₁₋₃;


each R' independently represents a member selected from the group consisting of: CONH₂; CONHC₁₋₆ alkyl(R^q)₃; CON(C₁₋₆ alkyl(R^q)₃)₂; CONHC₃₋₈ cycloalkyl(R^q)₃; CON(C₃₋₈ cycloalkyl(R^q)₃)₂; CON(C₃₋₈ cycloalkyl)(C₁₋₆ alkyl(R^q)₃); CO₂H; CO₂C₁₋₆ alkyl(R^q)₃; C(O)C₁₋₆ alkyl(R^q)₃; CO₂C₃₋₈ cycloalkyl(R^q)₃; C(O)C₃₋₈ cycloalkyl(R^q)₃; $-\text{[C(O)(CH}_2\text{)]}_j\text{-CR}^5\text{R}^6\text{-(CH}_2\text{)]}_k\text{-NR}^7\text{]}_p\text{-R}^8$; -C(O)C₃₋₈ cycloalkyl(R^q)₃; -C(O)heterocyclyl(R^q)₃; -CON[C₁₋₆ alkyl(R^q)₃][C₃₋₈ cycloalkyl(R^q)₃]; -C(O)aryl(R^q)₃;

-5-

- C(O)heteroaryl(R^q)₃; hydroxy; C₁₋₆ alkyl(R^q)₃; C₃₋₈ cycloalkyl(R^q)₃; OC₁₋₆ alkyl(R^q)₃; OC₃₋₈ cycloalkyl(R^q)₃; heterocyclyl(R^q)₃; CN; NH(R^{q'}); NHC₁₋₆ alkyl(R^q)₃; N(C₁₋₆ alkyl(R^q)₃)₂; NHC₃₋₈ cycloalkyl(R^q)₃; N(C₃₋₈ cycloalkyl(R^q)₃)₂;
 5 CF₃; SH; NO₂; C₂₋₄ alkenyl(R^q)₂₋₃, aryl(R^q)₃; heteroaryl(R^q)₃; C₂₋₄ alkynyl(R^q)₁₋₃; -OC(O) C₃₋₈ cycloalkyl(R^q)₃; SO₂NH₂; SO₂NHC₁₋₆ alkyl(R^q)₃; SO₂N(C₁₋₆ alkyl(R^q)₃)₂; NHSO₂C₁₋₆ alkyl(R^q)₃; NHSO₂aryl(R^q)₃; NHSO₂heteroaryl(R^q)₃;
 10 -OC(O)heterocyclyl(R^q)₃; N(R^{q'})C(O)C₁₋₆ alkyl(R^q)₃; NR^{q'}C(O)NH(C₁₋₆ alkyl(R^q)₃); -OC(O)C₁₋₆ alkyl(R^q)₃; -OC(O)aryl(R^q)₃; -OC(O)heteroaryl(R^q)₃; -C(=NR^{q'})NH₂; -C(=N^{q'})NHC₁₋₆ alkyl(R^q)₃; -C(=N^{q'})N(C₁₋₆ alkyl(R^q)₃)₂;



15



wherein j and k independently represent integers of from 0 -
 3;

20 R⁵ and R⁶ are independently H, aryl, C₁₋₆ alkyl(R^q)₃, or CR⁵R⁶ in combination represents a 3, 4, 5 or 6 membered cycloalkyl or heterocyclyl group, an aryl group or a heteroaryl group;


25 p represents 1, 2 or 3, with the proviso that when p represents 1, CR⁵R⁶ represents a 3, 4, 5 or 6 membered cycloalkyl group or a heterocyclyl group, an aryl group or a heteroaryl group, and at least one of j and k is 1, 2 or 3;



30 R⁷ and R⁸ are independently H, C₁₋₆ alkyl or aryl;

R⁹ represents H, a negative charge balanced by a positively charged group or a protecting group;

-6-

R^q represents a member selected from the group consisting of: R^{q'}; halo; CN; CO₂H; CO₂C₁₋₄ alkyl; C(O)C₁₋₄ alkyl; NH(R^{q''}); aryl(R^a)₃; heteroaryl(R^a)₃; NHC₁₋₄ alkyl; N(C₁₋₄ alkyl)₂; CONH₂; SH; S(O)_yC₁₋₆ alkyl(R^a)₃; C(O)NHC₁₋₆ alkyl(R^a)₃; C(O)N(C₁₋₆ alkyl(R^a)₃)₂; C₃₋₈ cycloalkyl; NHC(NH)NH₂; -heteroalkyl(R^a)₃; -NHC(O)NH₂;



wherein  and  independently represent mono or bicyclic ring systems, non-aromatic or partially aromatic, containing from 5-10 ring atoms, 1-4 of which are N and 0-1 of which are O or S(O)_y, with y equal to 0, 1 or 2, optionally containing 1-2 carbonyl groups;

each R^a independently represents a member selected from the group consisting of: H, C₁₋₆ alkyl, OC₁₋₆ alkyl, aralkyl, substituted aralkyl, heteroaralkyl, substituted heteroaralkyl, aralkoxy, substituted aralkoxy, halo, hydroxy, CN, CONH₂, CONHC₁₋₆ alkyl, CON(C₁₋₆ alkyl)₂, CO₂H, CO₂C₁₋₆ alkyl, C(O)C₁₋₆ alkyl, phenyl, CF₃, SH, NO₂, SO_yC₁₋₆ alkyl, with y as defined above; SO₂NH₂, SO₂NHC₁₋₆ alkyl, NHSO₂(substituted aryl), NHSO₂(substituted heteroaryl), NHSO₂C₁₋₆alkyl, NHSO₂aryl, NHSO₂heteroaryl, NH₂, NHC₁₋₆ alkyl, N(C₁₋₆ alkyl)₂, NHC(O)C₁₋₆ alkyl, NHC(O)NH(C₁₋₆ alkyl), C₂₋₄ alkenyl and C₂₋₄ alkynyl;

R^{q'} represents H, OH, C₁₋₄ alkyl, -OC₁₋₄ alkyl, aryl or C(O)C₁₋₄ alkyl, and

R^{q''} represents H, OH or OC₁₋₄ alkyl.

-7-

A pharmaceutical composition is also included in the invention described herein, which is comprised of a compound of formula I as defined above in combination with a pharmaceutically acceptable carrier.

5

Also included in the invention is a method of treating a cytokine mediated disease in a mammal, comprising administering to a mammalian patient in need of such treatment an amount of a compound of formula I which is effective for treating said cytokine mediated disease.

10

DETAILED DESCRIPTION OF THE INVENTION

The invention is described herein in detail using the terms defined below unless otherwise specified.

15


The term "alkyl" refers to a monovalent alkane (hydrocarbon) derived radical containing from 1 to 15 carbon atoms unless otherwise defined. It may be straight or branched, and when of sufficient size, e.g., C₃-15 may be cyclic. Preferred straight or branched alkyl groups include methyl, ethyl, propyl, isopropyl, butyl and t-butyl. Preferred cycloalkyl groups include cyclopropyl, cyclopentyl and cyclohexyl.

20

Alkyl also includes an alkyl group substituted with a cycloalkyl group, such as cyclopropylmethyl.

25

Alkyl also includes a straight or branched alkyl group which contains or is interrupted by a cycloalkylene portion. Examples include the following:



wherein: x' and y' = from 0-10; and w and z = from 0-9.

30

The alkylene and monovalent alkyl portion(s) of the alkyl group can be attached at any available point of attachment to the cycloalkylene portion.

-8-

When substituted alkyl is present, this refers to a straight, branched or cyclic alkyl group as defined above, substituted with 1-3 groups as defined with respect to each variable.

5 Heteroalkyl means an alkyl group containing from 2-15 carbon atoms and being interrupted by from 1-4 heteroatoms selected from O, S and N.

10 The term "alkenyl" refers to a hydrocarbon radical straight, branched or cyclic containing from 2 to 15 carbon atoms and at least one carbon to carbon double bond. Preferably one carbon to carbon double bond is present, and up to four non-aromatic (non-resonating) carbon-carbon double bonds may be present. Preferred alkenyl groups include ethenyl, propenyl, butenyl and cyclohexenyl. As described above with respect to alkyl, the straight, branched or cyclic portion of the alkenyl group may contain double bonds and may be substituted when a substituted alkenyl group is provided.

15 The term "alkynyl" refers to a hydrocarbon radical straight, branched or cyclic, containing from 2 to 15 carbon atoms and at least one carbon to carbon triple bond. Up to three carbon-carbon triple bonds may be present. Preferred alkynyl groups include ethynyl, propynyl and butynyl. As described above with respect to alkyl, the straight, branched or cyclic portion of the alkynyl group may contain triple bonds and may be substituted when a substituted alkynyl group is provided.

25 Aryl refers to aromatic rings e.g., phenyl, substituted phenyl and like groups as well as rings which are fused, e.g., naphthyl and the like. Aryl thus contains at least one ring having at least 6 atoms, with up to two such rings being present, containing up to 10 atoms therein, with alternating (resonating) double bonds between adjacent carbon atoms. The preferred aryl groups are phenyl and naphthyl. Aryl groups may likewise be substituted as defined below. Preferred substituted aryls include phenyl and naphthyl substituted with one or two groups.

30 The term "heteroaryl" refers to a monocyclic aromatic hydrocarbon group having 5 or 6 ring atoms, or a bicyclic aromatic

-9-

group having 8 to 10 atoms, containing at least one heteroatom, O, S or N, in which a carbon or nitrogen atom is the point of attachment, and in which one additional carbon atom is optionally replaced by a heteroatom selected from O or S, and in which from 1 to 3 additional carbon atoms are optionally replaced by nitrogen heteroatoms. The heteroaryl group is optionally substituted with up to three groups.

Heteroaryl includes aromatic and partially aromatic groups which contain one or more heteroatoms. Examples of this type are thiophene, purine, imidazopyridine, pyridine, oxazole, thiazole, oxazine, pyrazole, tetrazole, imidazole, pyridine, pyrimidine, pyrazine and triazine. Examples of partially aromatic groups are tetrahydroimidazo[4,5-c]pyridine, phthalidyl and saccharinyl, as defined below.

Each R^a independently represents a member selected from the group consisting of: H, C₁₋₆ alkyl, OC₁₋₆ alkyl, aralkyl, substituted aralkyl, heteroaralkyl, substituted heteroaralkyl, aralkoxy, substituted aralkoxy, halo, hydroxy, CN, CONH₂, CONHC₁₋₆ alkyl, CON(C₁₋₆ alkyl)₂, CO₂H, CO₂C₁₋₆ alkyl, C(O)C₁₋₆ alkyl, phenyl, CF₃, SH, NO₂, SO_yC₁₋₆ alkyl, with y as defined above; SO₂NH₂, SO₂NHC₁₋₆ alkyl, NHSO₂(substituted aryl), NHSO₂(substituted heteroaryl), NHSO₂C₁₋₆alkyl, NHSO₂aryl, NHSO₂heteroaryl, NH₂, NHC₁₋₆ alkyl, N(C₁₋₆ alkyl)₂, NHC(O)C₁₋₆ alkyl, NHC(O)NH(C₁₋₆ alkyl), C₂₋₄ alkenyl and C₂₋₄ alkynyl. In substituted aralkyl, substituted heteroaralkyl and substituted aralkoxy, the aryl, heteroaryl or alkyl portions thereof can be substituted as appropriate.

Substituted alkyl, aryl and heteroaryl, and the substituted portions of aralkyl, aralkoxy, heteroaralkyl, heteroaralkoxy and like groups are substituted with from 1-3 groups selected from the group consisting of: halo, hydroxy, cyano, acyl, acylamino, aralkoxy, alkylsulfonyl, arylsulfonyl, alkylsulfonylamino, arylsulfonylamino, alkylaminocarbonyl, alkyl, alkoxy, aryl, aryloxy, aralkoxy, amino, alkylamino, dialkylamino, and sulfonylamino.

The terms "heterocycloalkyl" and "heterocyclyl" refer to a cycloalkyl group (nonaromatic) in which one of the carbon atoms in the

-10-

ring is replaced by a heteroatom selected from O, S(O)_y or N, and in which up to three additional carbon atoms may be replaced by said heteroatoms. When three heteroatoms are present in the heterocycle, they are not all linked together.

5 Examples of heterocyclyls are piperidinyl, morpholinyl, azetidiny, pyrrolidinyl, tetrahydrofuranyl, imidazoliny, piperazinyl, pyrrolidin-2-one, piperidin-2-one and the like.

Acyl as used herein refers to -C(O)C₁₋₆ alkyl and -C(O)-aryl.

10 Acylamino refers to the group -NHC(O)C₁₋₆ alkyl and -NHC(O)aryl.

Aralkoxy refers to the group -OC₁₋₆ alkylaryl.

Alkylsulfonyl refers to the group -SO₂C₁₋₆ alkyl.

Alkylsulfonylamino refers to the group -NHSO₂C₁₋₆alkyl.

15 Arylsulfonylamino refers to the group -NHSO₂aryl.



Alkylaminocarbonyl refers to the group -C(O)NHC₁₋₆ alkyl.

Aryloxy refers to the group -O-aryl.



Sulfonylamino refers to the group -NHSO₃H.

Halo means Cl, F, Br and I selected on an independent basis.


20

 and  are optional substituents linked to the HETCy group.

25

 and  independently represent mono or bicyclic ring systems, non-aromatic or partially aromatic, containing from 5-10 ring atoms, 1-4 of which are N and 0-1 of which are O or S(O)_y, with y equal to 0, 1 or 2, and when partially aromatic, the non-aromatic portion thereof optionally containing 1-2 carbonyl groups. Hence, these ring


systems can be heteroaryl or heterocyclic as defined above.



-11-

is linked to HETCy through a nitrogen atom contained in the ring system, either directly or through a linking group which is part of R'. Examples

include phthalidyl and saccharinyl, as further defined below.




is

likewise linked to HETCy, but through a carbon atom contained in the ring system.


5

The term phthalidyl refers to the heteroaryl



group:


The term saccharinyl refers to the heteroaryl



10


group:

The group





15

means



20

and



-12-

means $-NR^Z(CH_2)_k-CR^5R^6-(CH_2)_j-C(O)-OR^9$,


$-NR^Z(CH_2)_k-CR^5R^6-(CH_2)_j-C(O)NR^7(CH_2)_kCR^5R^6(CH_2)_jCO_2R^9$

5 or

$-NR^7(CH_2)_kCR^5R^6(CH_2)_jC(O)NR^7(CH_2)_kCR^5R^6(CH_2)_jC(O)NR^7(CH_2)_kCR^5R^6(CH_2)_jCO_2R^9$

10 The variables are determined independently within each group. Thus, e.g., when more than one j is present, they may be the same or different. The values of R^5 and R^6 can be H, C₁₋₆ alkyl(R⁹)₃ or aryl, or CR^5R^6 taken in combination represents a 3-6 membered cycloalkyl or heterocyclyl group, an aryl group or a heteroaryl group. Examples of

15 suitable CR^5R^6 groups include:



20 The patterns of attachment noted above can also include heteroatoms as appropriate. When an aryl or heteroaryl group is represented by CR^5R^6 , attachment cannot be through the same carbon atom.

-13-

The term "TNF mediated disease or disease state" refers to disease states in which TNF plays a role, either by production or increased activity levels of TNF itself, or by causing another monokine to be released, such as but not limited to IL-1 or IL-6. A disease state in which IL-1, for instance is a major component, and whose production or action, is exacerbated or secreted in response to TNF, would therefore be considered a disease state mediated by TNF.

The term "cytokine" as used herein means any secreted polypeptide that affects the functions of cells and is a molecule which modulates interactions between cells in the immune, inflammatory or hematopoietic response. A cytokine includes, but is not limited to, monokines and lymphokines regardless of which cells produce them. Examples of cytokines include, but are not limited to, Interleukin-1 (IL-1), Interleukin-6 (IL-6), Interleukin-8 (IL-8), Tumor Necrosis Factor-alpha (TNF-a) and Tumor Necrosis Factor-beta (TNF-b).

By the term "cytokine interfering or cytokine suppressive amount" is mean an effective amount of a compound of formula I which will cause a decrease in the *in vivo* activity or level of the cytokine to normal or sub-normal levels, when given to the patient for the prophylaxis or therapeutic treatment of a disease state which is exacerbated by, or caused by, excessive or unregulated cytokine production or activity.

The compounds of the present invention may contain one or more asymmetric carbon atoms and may exist in racemic and optically active forms. All are within the scope of the present invention.

One subset of compounds of particular interest is described with respect to formula I wherein one or two R" groups are present, and each independently represents NH₂, NHC₁₋₆ alkyl(R^q)₃, N(C₁₋₆ alkyl)₂, NHC₃₋₈ cycloalkyl, N(R^q)C(O)C₁₋₆ alkyl(R^q)₃, C₁₋₆ alkyl(R^q)₃, OC₁₋₆ alkyl(R^q)₃, CO₂H, CONH₂, NR^qC(O)NHC₁₋₆ alkyl(R^q)₃ or heterocyclyl(R^q)₃. Within this subset of compounds, all other variables are as previously defined.

Another subset of interest includes compounds of formula I wherein HETCy represents a 5-6 membered non-aromatic



-14-

heterocycle with 1-2 nitrogen atoms contained therein. HETCy is preferably a pyrrolidinyl or piperidinyl group, and most preferably a 4-piperidinyl group. Within this subset of compounds, all other variables are as previously defined.

5 Another subset of compounds is represented by formula I wherein R' is selected from C₁₋₆ alkyl(R⁹)₃, OC₁₋₆ alkyl(R⁹)₃, -C(O)C₁₋₆ alkyl(R⁹)₃; CN, NO₂ and CO₂C₁₋₆ alkyl(R⁹)₃. Within this subset of compounds, all other variables are as previously defined.

10 Another subset of interest includes compounds of formula I wherein from 1-3 R groups are present and each independently represents a member selected from the group consisting of: halo, hydroxy, C₁₋₆ alkyl(R⁹)₃, OC₁₋₆ alkyl(R⁹)₃, NH₂, NHC₁₋₆ alkyl(R⁹)₃, N(C₁₋₆ alkyl(R⁹)₃)₂ and CF₃. Within this subset of compounds, all other variables are as previously defined.

15 More particularly, a subset of interest relates to compounds of formula I wherein one or two R groups are present, selected from halo and CF₃. Within this subset of compounds, all other variables are as previously defined.

20 Another subset of compounds is represented by formula I wherein R^x is H, C₃₋₈ cycloalkyl or C₁₋₆ alkyl(R⁹)₃. More particularly, R^x represents H, CH₃, CH₂CH₃, CH₂CH₂CH₃, CH₂CH₂CH₂CH₃,  or -CH₂-. Within this subset of compounds, all other variables are as previously defined.

25 Another subset of compounds is represented by formula I wherein X' represents a direct bond. Within this subset of compounds, all other variables are as previously defined.

30 Another subset of compounds is represented by formula I wherein X represents -(CH₂)_m-Y-(CH₂)_n-, Y represents a direct bond, O, S or C(O); m represents 0 or 1 and n represents 0 or 1. Within this subset, X preferably represents a direct bond. Within this subset of compounds, all other variables are as previously defined.

A preferred subset of compounds is represented by formula I wherein:

-15-

one or two R" groups are present, each independently representing NH₂, NHC₁₋₆ alkyl(R^q)₃, N(C₁₋₆ alkyl)₂, NHC₃₋₈ cycloalkyl, N(R^q)C(O)C₁₋₆ alkyl(R^q)₃, C₁₋₆ alkyl(R^q)₃, OC₁₋₆ alkyl(R^q)₃, CO₂H, CONH₂, NR^qC(O)NHC₁₋₆ alkyl(R^q)₃ or heterocyclyl(R^q)₃;

HETCy represents a 5-6 membered non-aromatic heterocyclyl with 1-2 nitrogen atoms contained therein;

R' is selected from C₁₋₆ alkyl(R^q)₃, OC₁₋₆ alkyl(R^q)₃, -C(O)C₁₋₆ alkyl(R^q)₃; CN; NO₂; and CO₂C₁₋₆ alkyl(R^q)₃ ;
 from 1-3 R groups are present and each independently represents a member selected from the group consisting of: halo, hydroxy, C₁₋₆ alkyl(R^q)₃, OC₁₋₆ alkyl(R^q)₃, NH₂, NHC₁₋₆ alkyl(R^q)₃, N(C₁₋₆ alkyl(R^q)₃)₂, CO₂H and CF₃ ;

R^x is H, C₃₋₈ cycloalkyl or C₁₋₆ alkyl(R^q)₃;

X' represents a direct bond;

X represents -(CH₂)_m-Y-(CH₂)_n-, Y represents a direct bond; m represents 0 or 1 and n represents 0 or 1.

A more preferred subset of compounds of the invention is represented by formula I wherein:

one or two R" groups are present, selected from the group consisting of: NH₂, NHC₁₋₆ alkyl(R^q)₃, N(C₁₋₆ alkyl)₂, C₁₋₆ alkyl(R^q)₃, OC₁₋₆ alkyl(R^q)₃ and heterocyclyl(R^q)₃;

HETCy represents a piperidinyl group;

zero or one R' is present, which is selected from C₁₋₆ alkyl(R^q)₃, -CO₂C₁₋₆ alkyl(R^q)₃ and -C(O)C₁₋₆ alkyl(R^q)₃;

one or two R groups are present, selected from halo and CF₃;


R^x is H, C₃₋₈ cycloalkyl or C₁₋₆ alkyl(R^q)₃; and

X and X' represent a direct bond.

A more preferred subset of compounds of the invention is represented by formula I wherein:

one R" group is present and is selected from:

-16-





5

HETCy represents a 4-piperidinyl group;

R' is absent;

one or two R groups are present selected from halo and




CF₃;

R^x represents H, CH₃  or $\text{-CH}_2\text{-}$  ;
 and X and X' represent direct bonds.


10

Representative species falling within the present invention include the following:

-17-





5




10

-18-





5





10







-19-









5









-20-



-21-




-22-



5

-23-



As used herein, THF means tetrahydrofuran, DMF means dimethylformamide and HOBT means hydroxybenzotriazole.

5 The compounds of the present invention are prepared by procedures illustrated in the accompanying schemes. The general method of preparing the imidazole nucleus is outlined in Scheme 1. 2-Mercaptopyrimidine 1 (commercially available from Aldrich Chemicals Inc) is reacted with a dialkyl dimethylformamide acetal such as dimethyl
10 formamide dimethyl acetal in solvent (e.g. toluene) in the presence of a base e.g. diisopropylethylamine. Deprotonation of the product 2 with a strong base such as lithium diisopropylamide and quenching of the anion with an appropriately substituted N,O-dimethylhydroxamide 3 provides a
15 ketone 4. Formation of the oximoketone 5 may be carried out under acidic conditions with either an inorganic nitrite (e.g. sodium nitrite) or an organic nitrite (e.g. t-butyl nitrite). Reaction of the oximinoketone with a suitably functionalized and protected amino aldehyde and an ammonium salt (e.g. ammonium acetate) in acetic acid leads to a
20 hydroxyimidazole 6. Reduction to the imidazole 7 may be accomplished by a number of methods described in the literature (hydrogenation, reaction with PCl_3 or titanium trichloride). The imidazole may then be alkylated either by reaction with an alkyl halide and base in solvent (e.g. methyl iodide, cesium carbonate, DMF) or by heating with a dimethylformamide dialkyl acetal neat or in a solvent (e.g.

-24-

dimethylformamide dimethyl acetal in toluene) to provide a mixture of isomers **8** and **9**, separable by chromatography.

As shown in Scheme 2, the methylthiopyrimidine can be oxidized to the methylsulfonylpyrimidine, for example with oxone in methanol and then the methylsulfonyl group displaced with an amine to provide an aminopyrimidine derivative. Deprotection of the benzyloxycarbonyl protecting group could then be accomplished by treatment with hydrogen bromide in acetic acid or aqueous hydrochloric acid.

Scheme 3 illustrates compounds that may be obtained after the amine displacement reaction described in Scheme 2, where the amine utilized was 4-methoxybenzylamine. Reduction of the intermediate **10** with lithium aluminum hydride affords the N-methypiperidine derivative **13**. Treatment with hydrogen bromide in acetic acid provides the piperidine **12** and treatment with hydrochloric acid at reflux leads to **11**. The piperidine of compound **12** can be further reacted for example by reductive alkylation to provide **14** or by acylation to provide compounds **15**.


Alternatively the methanesulfonyl group can be displaced with cyanide as shown in Scheme 4 to afford **17**. Compound **17** may then be converted to a number of different products. Utilizing refluxing hydrochloric acid leads to the parent pyrimidine **19**, whereas hydrogen bromide in acetic acid gives the carboxamide **20**, which can be hydrolyzed to the acid **21** under basic conditions, e.g. aqueous sodium hydroxide solution.

Reactions on the piperidine nitrogen may also be carried out prior to the displacement of the sulfonylpyrimidine as shown in Scheme 5. The piperidine nitrogen of **22** may be acylated, for example employing EDC, HOBt and triethylamine and an appropriate carboxylic acid in DMF to give **23** and then the sulfonyl group displaced with an amine, e.g. ammonia to provide **24**.

-25-


The aminopyrimidine can also be acylated, for example by treatment with an appropriate acid chloride (Scheme 6) to give **26** from which the protecting group can be removed with hydrogen bromide in acetic acid, yielding compounds **27**.

5



SCHEME 1

-26-


SCHEME 1 cont'd




alkyl halide/base
or
dimethylformamide
dialkyl acetal
↓
heat




-27-

SCHEME 2


-28-

Scheme 2 cont'd


-29-

SCHEME 3

-30-


Scheme 3 cont'd

-31-


Scheme 4

-32-




Scheme 4 cont'd



-33-

Scheme 5

-34-

Scheme 6

The compounds of the present invention are useful in various pharmaceutically acceptable salt forms. The term

-35-

"pharmaceutically acceptable salt" refers to those salt forms which would be apparent to the pharmaceutical chemist. i.e., those which are substantially non-toxic and which provide the desired pharmacokinetic properties, palatability, absorption, distribution, metabolism or excretion.

5 Other factors, more practical in nature, which are also important in the selection, are cost of the raw materials, ease of crystallization, yield, stability, hygroscopicity and flowability of the resulting bulk drug. Conveniently, pharmaceutical compositions may be prepared from the active ingredients in combination with pharmaceutically acceptable
10 carriers.

The pharmaceutically acceptable salts of the compounds of formula I include conventional non-toxic salts or quarternary ammonium salts of the compounds of formula I formed e.g. from non-toxic inorganic or organic acids. For example, non-toxic salts include
15 those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methane-
20 sulfonic, ethane disulfonic, oxalic, isethionic, trifluoroacetic and the like.

The pharmaceutically acceptable salts of the present invention can be synthesized by conventional chemical methods. Generally, the salts are prepared by reacting the free base or acid with
25 stoichiometric amounts or with an excess of the desired salt-forming inorganic or organic acid or base, in a suitable solvent or solvent combination.

The compounds of the present invention may have asymmetric centers and occur as racemates, racemic mixtures, and as
30 individual diastereomers. All such isomers, including optical isomers, being included in the present invention.

The compounds of formula I can be used in the prophylactic or therapeutic treatment of disease states in mammals which are

-36-

exacerbated or caused by excessive or unregulated cytokines, e.g., IL-1, IL-6, IL-8 or TNF.

5 Because the compounds of formula I inhibit cytokines, the compounds are useful for treating diseases in which cytokine presence or activity is implicated, such as rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions.

10 The compounds of formula I are useful to treat disease states mediated by excessive or unregulated TNF production or activity. Such diseases include, but are not limited to sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, adult respiratory distress syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcoidosis, bone resorption diseases, such as osteoporosis, reperfusion injury, graft v. host rejection, allograft rejection, fever, myalgia due to infection, cachexia secondary to infection or malignancy, cachexia secondary to acquired immune deficiency syndrome (AIDS), AIDS, ARC (AIDs related complex), keloid formation, scar tissue formation, Crohn's disease, ulcerative colitis, pyresis, AIDS and other viral infections, such as cytomegalovirus (CMV), influenza virus, and the herpes family of viruses such as Herpes 20 Zoster or Simplex I and II.

25 The compounds of formula I are also useful topically in the treatment of inflammation such as in the treatment of rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions; inflamed joints, eczema, psoriasis or other inflammatory skin conditions such as sunburn; inflammatory eye conditions including conjunctivitis; pyresis, pain and other conditions associated with inflammation.

30 The compounds of formula I are also useful in treating diseases characterized by excessive IL-8 activity. These disease states include psoriasis, inflammatory bowel disease, asthma, cardiac and renal reperfusion injury, adult respiratory distress syndrome, thrombosis and glomerulonephritis.

The invention thus includes a method of treating psoriasis, inflammatory bowel disease, asthma, cardiac and renal reperfusion injury,

-37-

adult respiratory distress syndrome, thrombosis and glomerulonephritis, in a mammal in need of such treatment, which comprises administering to said mammal a compound of formula I in an amount which is effective for treating said disease or condition.

5 When administered to a patient for the treatment of a disease in which a cytokine or cytokines are implicated, the dosage used can be varied within wide limits, depending upon the type of disease, the age and general condition of the patient, the particular compound administered, the presence or level of toxicity or adverse effects experienced with the
10 drug and other factors. A representative example of a suitable dosage range is from as low as about 0.01 mg/kg to as high as about 100 mg/kg. However, the dosage administered is generally left to the discretion of the physician.

 The methods of treatment can be carried out by delivering
15 the compound of formula I parenterally. The term 'parenteral' as used herein includes intravenous, intramuscular, or intraperitoneal administration. The subcutaneous and intramuscular forms of parenteral administration are generally preferred. The instant invention can also be carried out by delivering the compound of formula I subcutaneously,
20 intranasally, intrarectally, transdermally or intravaginally.

 The compounds of formula I may also be administered by inhalation. By 'inhalation' is meant intranasal and oral inhalation administration. Appropriate dosage forms for such administration, such as an aerosol formulation or a metered dose inhaler, may be prepared by
25 convention techniques.

 The invention also relates to a pharmaceutical composition comprising a compound of formula I and a pharmaceutically acceptable carrier. The compounds of formula I may also be included in
30 pharmaceutical compositions in combination with a second therapeutically active compound.

 The pharmaceutical carrier employed may be, for example, either a solid, liquid or gas. Examples of solid carriers include lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Examples of liquid carriers are syrup, peanut

-38-

oil, olive oil, water and the like. Examples of gaseous carriers include carbon dioxide and nitrogen.

Similarly, the carrier or diluent may include time delay material well known in the art, such as glyceryl monostearate or glyceryl distearate, alone or with a wax.

A wide variety of pharmaceutical dosage forms can be employed. If a solid dosage is used for oral administration, the preparation can be in the form of a tablet, hard gelatin capsule, troche or lozenge. The amount of solid carrier will vary widely, but generally will be from about 0.025 mg to about 1 g. When a liquid dosage form is desired for oral administration, the preparation is typically in the form of a syrup, emulsion, soft gelatin capsule, suspension or solution. When a parenteral dosage form is to be employed, the drug may be in solid or liquid form, and may be formulated for administration directly or may be suitable for reconstitution.

Topical dosage forms are also included. Examples of topical dosage forms are solids, liquids and semi-solids. Solids would include dusting powders, poultices and the like. Liquids include solutions, suspensions and emulsions. Semi-solids include creams, ointments, gels and the like.

The amount of a compound of formula I used topically will, of course, vary with the compound chosen, the nature and severity of the condition, and can be varied in accordance with the discretion of the physician. A representative, topical, dose of a compound of formula I is from as low as about 0.01 mg to as high as about 2.0 g, administered one to four, preferably one to two times daily.

The active ingredient may comprise, for topical administration, from about 0.001 % to about 10% w/w.

Drops according to the present invention may comprise sterile or non-sterile aqueous or oil solutions or suspensions, and may be prepared by dissolving the active ingredient in a suitable aqueous solution, optionally including a bactericidal and/or fungicidal agent and/or any other suitable preservative, and optionally including a surface active agent. The resulting solution may then be clarified by filtration,

-39-

transferred to a suitable container which is then sealed and sterilized by autoclaving or maintaining at 98-100°C for half an hour. Alternatively, the solution may be sterilized by filtration and transferred to the container aseptically. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily solution include glycerol, diluted alcohol and propylene glycol.


Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those for the preparation of drops. Lotions or liniments for application to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

Creams, ointments or pastes according to the present invention are semi-solid formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely-divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous liquid, with a greasy or non-greasy base. The base may comprise hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage; an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives, or a fatty acid such as stearic or oleic acid together with an alcohol such as propylene glycol or macrogels. The formulation may incorporate any suitable surface active agent such as an anionic, cationic or non-ionic surfactant such as sorbitan esters or polyoxyethylene derivatives thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as silicas, and other ingredients such as lanolin may also be included.

-40-

EXAMPLE 14-METHOXYBENZYL-[4-[2-PIPERIDIN-4-YL-5-(3-TRIFLUOROMETHYLPHENYL)-3H-IMIDAZOL-4-YL]-PYRIMIDIN-2-YL]AMINE

5

Step 1-A2-Methylthio-4-methylpyrimidine

10

To 2-mercapto-4-methylpyrimidine•HCl (50.0g, 0.307mole) in toluene (750 mL), under argon, was added diisopropylethylamine (80.0 mL, 0.461 mole) followed by N,N-dimethylformamide dimethyl acetal (100 mL) and the mixture heated to reflux for 4 hours. Upon cooling, the reaction was concentrated *in vacuo* to an oil, dissolved in ether (400 mL), and washed with water (2x50 mL). The organic extract was dried over anhydrous sodium sulfate, filtered and concentrated to an oil which was vacuum distilled to give 2-methylthio-4-methylpyrimidine (36.4 g) as an oil.

20

¹H NMR(CDCl₃) δ 8.37 (d, 1H, J = 7.5Hz), 6.82 (d, 1H, J = 7.5Hz), 2.55 (s 3H), 2.45 (s, 3H).

-41-

Step 1-B2-(2-Methylthiopyrimidin-4-yl)-1-(3-trifluoromethylphenyl)ethanone

To a solution of diisopropylamine (7.9 mL, 0.056 mole) in THF (100 mL) at -78°C under argon was added 2.5M N-butyllithium (22.5 mL, 0.056 mole), followed after 5 minutes by a solution of 2-methylthio-4-methylpyrimidine (5.27 g, 0.0376 mole) in THF (20 mL). Upon stirring for 15 min. at -78°C, a solution of N-methoxy-N-methyl-3-trifluoromethylbenzamide (9.63 g, 0.041 mole) in THF (90 mL) was added. The reaction was allowed to warm to 0°C and then quenched by pouring into water (400 mL) and ethyl acetate (400 mL). The layers were separated and the aqueous layer washed with ethyl acetate (200 mL). The ethyl acetate extracts were combined, dried over anhydrous sodium sulfate, filtered, and concentrated to a solid (11.9 g). Trituration with 10% ether/hexane (100 mL) gave 9.5 g of the title compound.

¹H NMR (CDCl₃) (mixture of keto-and enol tautomers) δ 6.0-8.5 (m, 6H, rotamers), 2.4-2.7 (m, 3H), .

Step 1-C1-(2-Methylthiopyrimidin-4-yl)-2-(3-trifluoromethylphenyl)-ethane-1,2-dione 1-oxime

To a mixture of 2-(2-methylthiopyrimidin-4-yl)-1-(3-trifluoromethylphenyl)ethanone (4.5 g, 0.0144 mole) in acetic acid (67 mL), was added THF (54 mL) and water (9 mL). The mixture was cooled to +5°C and a solution of sodium nitrite (1.34 g, 0.0194 mole) added dropwise while maintaining the temperature below +10°C. Upon completion of addition, the reaction was allowed to warm to room temperature for 1 hour, diluted with water (200 mL) and ethyl acetate (200 mL) and the pH adjusted to 7.5 with 3N NaOH. The layers were separated and the aqueous layer washed with ethyl acetate (100 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated to an oil. Yield 4.9 g.

-42-

¹H NMR (CDCl₃) δ 8.58 (d, 1H, J = 5.4 Hz), 8.18 (s, 1H), 8.06 (d, 1H, J = 8.6 Hz), 7.89 (d, 1H, J = 8.6 Hz), 7.66 (t, 1H, J = 8.6 Hz), 7.52 (d, 1H, J = 5.4 Hz), 2.18 (s, 3H), 1.60 (brs, 1H).

5 Step 1-D

4-[1-Hydroxy-5-(2-methylthiopyrimidin-4-yl)-4-(3-
 trifluoromethylphenyl)-1H-imidazol-2-yl]piperidine-1-carboxylic acid
 benzyl ester

 To a mixture of 1-(2-methylthiopyrimidin-4-yl)-2-(3-
10 trifluoromethylphenyl)ethane-1,2-dione-1-oxime (5.0 g, 0.0146 mole)
 and N-carbobenzyloxypiperidine-4-carboxaldehyde [Amici *et al* Eur. J.
 Med. Chem. **26**, 625-631 (1991)] (4.70 g, 0.019 mole), in acetic acid (140
 mL) was added ammonium acetate (23 g, 0.295 mole). The mixture was
15 heated to reflux for 1-1/2 hours, cooled and concentrated to remove most
 of the acetic acid. The residue was dissolved in water (200 mL) and ethyl
 acetate (400 mL) and the pH adjusted to 7.5 with 3N sodium hydroxide.
 The ethyl acetate layer was removed, dried over Na₂SO₄, filtered and
 concentrated to give an oil (9.5 g crude product). The oil was used in the
20 next step without further purification.

Step 1-E

4-[5-(2-Methylthiopyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-
 imidazol-2-yl]piperidine-1-carboxylic acid benzyl ester

 To a stirring solution of 4-[1-hydroxy-5-(2-
25 methylthiopyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-
 yl]piperidine-1-carboxylic acid benzyl ester (9.5 g, crude, 0.0146 mole)
 in methanol (130 mL) at 20°C was added titanium (III) chloride (25 mL,
 0.029 mole, 15% wt in 20-30% HCl) dropwise over 10 minutes. The
 reaction was allowed to stir for 3 hours then quenched by pouring slowly
30 into a mixture of 10% aqueous sodium bicarbonate (1.5 L) and ethyl
 acetate (600 mL). After stirring for 30 minutes the organic layer was
 removed and aqueous extract washed with ethyl acetate (2x200 mL). The
 organic extracts were combined, dried over anhydrous sodium sulfate,
 and concentrated to an oil. The oil was chromatographed on silica using

-43-

60% ethyl acetate/hexane to give 6.35 g of a yellow foam upon concentration of product containing fractions.

¹H NMR (CDCl₃) δ 10.18 (brs, 1H), 8.28 (d, 1H, J = 5.5 Hz), 7.2-7.8 (m, 9H), 6.95 (d, 1H, J = 5.5 Hz), 5.12 (s, 2H), 4.30 (brs, 2H), 3.0 (brm, 3H), 2.58 (s, 3H), 1.7-2.2 (m, 4H).

Step 1-F

4-[5-(2-Methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]piperidine-1-carboxylic acid benzyl ester

To a stirring solution of 4-[5-(2-methylthiopyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]piperidine-1-carboxylic acid benzyl ester (2.5 g, 0.0045 mole) in methanol (75 mL) at 20°C was slowly added an aqueous solution (75 mL) of Oxone® (8.32 g, 0.0135 mole). The reaction was allowed to stir for 4 hours, concentrated *in vacuo* to remove methanol, diluted with 10% aqueous sodium bicarbonate (100 mL), and extracted with ethyl acetate (2x150 mL). The organic extracts were combined, dried over anhydrous sodium sulfate, filtered and concentrated to give the title compound (2.75 g).

¹H NMR (CDCl₃) δ 11.06 (s, 1H), 7.5-8.6 (m, 6H), 7.35 (m, 5H), 5.12 (s, 2H), 4.30 (brs, 2H), 3.35 (s, 3H), 2.7-3.1 (m, 3H), 1.8-2.2 (m, 4H).

Step 1-G

4-[5-(2-(4-Methoxybenzylamino)-pyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester

A mixture of 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (1.5 g, 0.00256 mole) and 4-methoxybenzylamine (3.51 g, 0.026 mole) was heated in a pressure tube at 140°C for 10 minutes. The mixture was allowed to cool and the residue column chromatographed on silica using 5% methanol/methylene chloride to give 1.52 g of a yellow powder upon concentration *in vacuo* of product containing fractions.

-44-

Anal. Calc'd for C₃₅H₃₃N₆O₃F₃: C 65.41, H 5.18, N 13.08.

Found: C 65.12, H 5.29, N 12.98.

Step 1-H

5 4-Methoxybenzyl-[4-[2-piperidin-4-yl]-5-(3-trifluoromethylphenyl)-3H-imidazol-4-yl]-pyrimidin-2-yl]amine

 To a stirred solution of 4-[5-[2-(4-methoxybenzylamino)-pyrimidin-4-yl]-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (1.0 g, 0.00156 mole) in
10 methylene chloride (16 mL) under argon was slowly added 30% hydrogen bromide in acetic acid (16 mL). The mixture was stirred at 20°C for 30 minutes and then diluted with diethyl ether (160 mL). The resulting mixture was stirred for 1 hour, filtered and solid washed with ether (10 mL). The resulting solid was dissolved in 10% aqueous sodium
15 bicarbonate (40 mL) and methylene chloride (50 mL). The methylene chloride was separated and the aqueous layer washed with methylene chloride (25 mL). The organic extracts were dried over anhydrous sodium sulfate, filtered and concentrated to a foam (0.80 g). The solid was chromatographed on silica using methylene
20 chloride/methanol/aqueous ammonium hydroxide (90:10:2) to give upon concentration of product containing fractions 180 mg of the title compound.


25 ¹H NMR (CDCl₃) δ 9.8 (brs, 1H), 8.12 (d, 1H, J = 5.8 Hz), 7.90 (s, 1H), 7.80 (d, 1H, J = 8.1 Hz), 7.62 (d, 1H, J = 8.1 Hz), 7.42 (t, 1H, J = 8.1 Hz), 7.32 (d, 1H, J = 8.1 Hz), 6.90 (d, 2H, J = 8.1 Hz), 6.78 (brs, 1H), 5.44 (brs, 1H), 4.6 (brs, 2H), 3.80 (s, 3H), 3.22 (m, 2H), 2.95 (m, 1H), 2.77 (m, 2H), 1.6-2.2 (m, 4H).

30 The following examples (2-17) were prepared as described in Example 1 and were isolated either by silica gel chromatography to give the free base, preparative HPLC to give the trifluoroacetic acid salt upon lyophilization, or crystallization of the hydrobromide salts.

-45-

EXAMPLE 24-[5-(2-METHYLAMINOPYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINE-1-CARBOXYLIC ACID BENZYL ESTER

5




To a stirred solution of 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F) (0.5 g, 0.8 mmole) in ethanol (10 mL) in a pressure vessel (50 mL) was added 40% aqueous methylamine (20 mL). The vessel was sealed and then heated at 100°C for 3 hours. The reaction was cooled, concentrated *in vacuo* and chromatographed on silica using methylene chloride/methanol/aqueous ammonium hydroxide (95:5:1) to give 0.42 g of product. Crystallization from ether (15 mL) gave 0.30 g of the title compound as a white solid.

Anal. Calc'd for C₂₈H₂₇N₆O₂F₃: C 62.68, H 5.07, N 15.66.

Found: C 62.89, H 5.05, N 15.92.

20

-46-

EXAMPLE 34-[5-(2-METHYLAMINOPYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINE


5

The title compound was prepared from 4-[5-[2-(methylamino)-pyrimidin-4-yl]-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F as
10 described in Example 1 step H.

Anal Calc'd for C₂₀H₂₁N₆F₃: C 59.69, H 5.26, N 20.88

Found: C 59.62, H 5.31, N 20.95

-47-

EXAMPLE 4**4-[5-(2-DIMETHYLAMINOPYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINE**

5


The title compound was prepared in a similar manner from 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F) as described in Example 1 steps G (replacing 4-methoxybenzylamine with 40% aqueous dimethylamine) and H.

¹H NMR (CDCl₃) δ 9.9 (brs, 1H), 8.15 (d, 1H, J = 5.0 Hz), 7.9 (s, 1H), 7.81 (d, 1H, J = 7.4 Hz), 7.62 (d, 1H, J = 7.4 Hz), 7.54 (t, 1H, J = 7.4 Hz), 6.54 (brs, 1H), 3.2 (m, 6H), 3.0 (m, 1H), 2.78 (m, 2H), 1.7-2.2 (m, 6H).

15

Mass Spectral Analysis - M⁺ = 417.

-48-

EXAMPLE 54-[5-[2-(1-PIPERDINYL)-PYRIMIDIN-4-YL]-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINE

5

The title compound was prepared from 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F) as described in Example 1 steps G (replacing 4-methoxybenzylamine with piperidine) and H.

¹H NMR (CDCl₃) δ 9.8 (brs, 1H), 8.15 (d, 1H, J = 5.5 Hz), 7.90 (s, 1H), 7.80 (d, 1H, J = 7.8 Hz), 7.62 (d, 1H, J = 7.8 Hz), 7.52 (t, 1H, J = 7.8 Hz), 3.80 (brs, 4H), 3.22 (m, 2H), 3.0 (m, 1H), 2.77 (m, 2H), 1.4-2.2 (m, 10H).


15

-49-

EXAMPLE 6

(R)-4-[5-(2-(1-PHENYLETHYLAMINO)-PYRIMIDIN-4-YL)-4-(3-
TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-
PIPERIDINE-1-CARBOXYLIC ACID BENZYL ESTER

5



10 The title compound was prepared from 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F) as described in Example 1 steps (replacing 4-methoxybenzylamine with R(+)-a-methylbenzylamine).


15 Anal. Calc'd for $C_{35}H_{33}N_6O_2F_3 \cdot 0.3H_2O$: C 66.50, H 5.36, N 13.30
Found: C 66.52, H 5.27, N 13.32

MP 85-87°C

-50-

EXAMPLE 7(R)-4-[5-(2-(1-PHENYLETHYLAMINO)-PYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINE
HYDROCHLORIDE

5



10 The title compound was prepared from (R)-4-[5-(2-(1-phenylethylamino)-pyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 6) as described in Example 1 step H.


15 MP = 155-160°C
[α]_D = +165.1°(MeOH)

-51-

EXAMPLE 8

(S)-4-[5-[2-(1-PHENYLETHYLAMINO)-PYRIMIDIN-4-YL]-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-
PIPERIDINE-1-CARBOXYLIC ACID BENZYL ESTER

5



The title compound was prepared from 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F) as described in Example 1 step G (replacing 4-methoxybenzylamine with S(-)-a-methylbenzylamine).


15 Anal. Calc'd for $C_{35}H_{33}N_6O_2F_3 \cdot 0.2H_2O$: C 66.69, H 5.34, N 13.33
Found: C 66.74, H 5.38, N 12.98

MP 85-87°C.

-52-

EXAMPLE 9

(S)-4-[5-(2-(1-PHENYLETHYLAMINO)-PYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINE



10 The title compound was prepared from (S)-4-[5-[2-(1-phenylethylamino)-pyrimidin-4-yl]-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 8) as described in Example 1 step H.

MP 157-161°C.


[α]_D = -165.1°(MeOH)

-53-

EXAMPLE 10

4-[5-(2-(CYCLOPROPYLAMINO)PYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINE-1-CARBOXYLIC ACID BENZYL ESTER

5




10 The title compound was prepared from 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F) as described in Example 1 step G (replacing 4-methoxybenzylamine with cyclopropylamine).

15 Mass Spectral Analysis = $M^{+1} = 563$.

-54-

EXAMPLE 11

4-[5-(2-(CYCLOPROPYLAMINO)PYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINE



5

The title compound was prepared 4-[5-(2-(Cyclopropylamino)pyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 10) as described in Example 1 step H.


¹H NMR(CDCl₃) δ 10.1 (brs, 1H), 8.16 (d, 1H, J = 5.3 Hz), 7.92 (s, 1H), 7.80 (d, 1H, J = 8.0 Hz), 7.62 (d, 1H, J = 8.0 Hz), 7.54 (t, 1H, J = 8.0 Hz), 6.62 (brs, 1H), 3.20 (m, 2H), 3.0 (m, 1H), 2.75 (m, 2H), 1.7-2.2 (m, 5H), 0.5-1.0 (m, 4H).

15

-55-

EXAMPLE 124-[5-(2-AMINOPYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINE-1-CARBOXYLIC ACID BENZYL ESTER

5



To a 50 mL pressure vessel cooled to -50°C and containing 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F) (100 mg, 0.176 mmole) was introduced liquid ammonia (10 mL). The vessel was sealed, allowed to warm to 25°C and stirred for 18 hours. The ammonia was allowed to evaporate and the residue purified by preparative HPLC using a C-18 column and 0.1% trifluoroacetic acid/water and acetonitrile as eluants. The title compound was isolated from the product containing fractions by lyophilization to yield 100 mg.

Mass Spectral Analysis - $M+1 = 523$


Anal. Calc'd for $\text{C}_{27}\text{H}_{25}\text{N}_6\text{O}_2\text{F}_3 \cdot 2.0\text{CF}_3\text{COOH}$: C 49.60, H 3.63, N 11.20

Found: C 49.53, H 3.40, N 10.90

-56-

EXAMPLE 13

4-[5-(2-(2,2,2-TRIFLUOROETHYLAMINO)-PYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINE



5


The title compound was prepared from 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F) as described in Example 1
10 steps G (replacing 4-methoxybenzylamine with 2,2,2-trifluoromethylamine and carrying out the reaction for 7 days at 140°C) and H.

15 Mass Spectral Analysis - $M^+1 = 471$
MP - 205-210°C

-57-

EXAMPLE 144-TRIFLUOROMETHYLBENZYL-[4-[2-PIPERIDIN-4-YL-5-(3-TRIFLUOROMETHYLPHENYL)-3H-IMIDAZOL-4-YL]-PYRIMIDIN-2-YL] AMINE

5



The title compound was prepared from 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F) as described in Example 1 steps G (replacing 4-methoxybenzylamine with 4-trifluoromethylbenzylamine) and H.

Purification by preparative HPLC resulted in isolation of the trifluoroacetic acid salt after lyophilization of product containing fractions.

Anal Calc'd for $C_{27}H_{24}N_6F_6 \cdot 2.8CF_3COOH$: C 45.22, H 3.12, N 9.71
Found: C 45.48, H 3.22, N 9.83


MP 75-79°C

-58-

EXAMPLE 15

3-TRIFLUOROMETHYLBENZYL-[4-[2-PIPERIDIN-4-YL-5-(3-TRIFLUOROMETHYLPHENYL)-3H-IMIDAZOL-4-YL]-PYRIMIDIN-2-YL] AMINE

5



The title compound was prepared from 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F) as described in Example 1 steps G (replacing 4-methoxybenzylamine with 3-trifluoromethylbenzylamine) and H.

15 Purification by preparative HPLC resulted in isolation of the trifluoroacetic acid salt after lyophilization of product containing fractions.

Anal. Calc'd for $C_{27}H_{24}N_6F_6 \cdot 2.8CF_3COOH$: C 45.22, H 3.12, N 9.71
Found: C 45.31, H 3.24, N 9.71


20 MP 68-74°C

-59-

EXAMPLE 16

2-TRIFLUOROMETHYLBENZYL-[4-[2-PIPERIDIN-4-YL-5-(3-TRIFLUOROMETHYLPHENYL)-3H-IMIDAZOL-4-YL]-PYRIMIDIN-2-YL] AMINE

5




The title compound was prepared from 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F) as described in Example 1 steps G (replacing 4-methoxybenzylamine with 2-trifluoromethylbenzylamine) and H.

15 Purification by preparative HPLC resulted in isolation of the trifluoroacetic acid salt after lyophilization of product containing fractions.

Anal. Calc'd for $C_{27}H_{24}N_6F_6 \cdot 3.0CF_3COOH$: C 44.15, H 3.14, N 9.36
Found: C 44.16, H 3.15, N 9.21

20 MP 65-68°C

-60-

EXAMPLE 174-[5-(2-(2-INDANYLAMINO)-PYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINE

5

The title compound was prepared from 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F) as described in Example 1 steps G (replacing 4-methoxybenzylamine with 2-aminoindane) and H.

The product was isolated as the trihydrobromide salt.

Anal. Calc'd for C₂₈H₂₇N₆F₃·3HBr, 1.5 H₂O: C 43.41, H 4.29, N 10.85


Found: C 43.78, H 4.59, N 10.49

-61-

EXAMPLE 18

[4-[2-(1-METHYL-PIPERIDIN-4-YL)-5-(3-TRIFLUOROMETHYLPHENYL)-3H-IMIDAZOL-4-YL]-PYRIMIDIN-2-YL]-4-METHOXYBENZYLAMINE

5



To a stirred solution of 4-[5-(2-(4-methoxybenzylamino)-pyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1G) (0.227 g, 0.000353 mole) in THF (10 mL) was slowly added 1.0M lithium aluminum hydride in THF (0.53 mL, 0.0053 mole). The reaction was warmed to reflux for 30 min., cooled and water (200 mL) added dropwise. The reaction mixture was diluted with THF (20 mL) and sodium sulfate (1.0 g) added. The mixture was stirred well for 10 minutes, filtered, concentrated *in vacuo*, and chromatographed on silica using methylene chloride/methanol/aqueous ammonium hydroxide (90:10:2) to give 100 mg of the title compound upon concentration of product containing fractions. The residue was triturated from 10% ether/hexane to give 70 mg of the title compound as a white solid.

¹H NMR (CDCl₃) δ 9.85 (brs, 1H), 8.10 (brd, 1H), 7.91 (s, 1H), 7.80 (d, 1H, J = 8.1), 7.5-7.7 (m, 2H), 7.31 (d, 2H, J = 8.1 Hz), 6.90 (d, 2H, J = 8.1), 6.59 (brd, 1H), 5.45 (brs, 1H), 4.58 (d, 2H, J = 5.6Hz), 3.80 (s, 3H), 2.8-3.0 (m, 3H), 2.32 (s, 3H), 1.8-2.2 (m, 6H).

-62-


Anal Calc'd for $C_{28}H_{29}N_6OF_3 \cdot 0.3 H_2O$: C 63.69, H 5.65, N 15.92

Found: C 63.70, H 5.70, N 15.94

5

EXAMPLE 19

1-[4-[5-(2-(4-METHOXYBENZYLAMINO)-PYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINE-YL]-ETHANONE



10

To a stirred solution of 4-methoxybenzyl-[4-[2-piperidin-4-yl-5-(3-trifluoromethylphenyl)-3H-imidazol-4-yl]-pyrimidin-2-yl]amine (Example 1H) (0.150 g, 0.295 mmole) in DMF (4 mL) was added acetic acid (18 mL, 0.31 mmole), triethylamine (62 mL, 0.44 mmole), 1-hydroxybenzotriazole hydrate (68 mg, 0.44 mmole) and EDC (85 mg, 0.44 mmole). The mixture was stirred at 20°C for 18 hours then diluted with 10% aqueous sodium bicarbonate (20 mL) and ethyl acetate (30 mL). The organic extract was removed, washed with water (10 mL), dried over anhydrous sodium sulfate, filtered, and column chromatographed on silica using methylene chloride/methanol/aqueous ammonium hydroxide (95:5:1) to give 146 mg. Crystallization from ether gave 120 mg of the title compound.

20

Anal Calc'd for $C_{29}H_{29}N_6O_2F_3$: C 63.20, H 5.31, N 15.26

25


Found: C 62.80, H 5.29, N 15.16

-63-

EXAMPLE 20

1-[4-[5-(2-(4-METHOXYBENZYLAMINO)PYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINYL]-2-DIMETHYLAMINOETHANONE

5



The following compound was prepared from 4-methoxybenzyl-[4-[2-piperidin-4-yl-5-(3-trifluoromethylphenyl)-3H-imidazol-4-yl]-pyrimidin-2-yl]amine (Example 1H) as described in example 19 using N'-N'-dimethylglycine in place of acetic acid to give the title compound.

¹H NMR (CDCl₃) δ 9.78 (brs, 1H), 8.12 (d, 1H, J = 5.8 Hz), 7.90 (s, 1H), 7.80 (d, 1H, J = 8.1 Hz), 7.64 (d, 1H, J = 8.1 Hz), 7.54 (t, 1H, J = 8.1 Hz), 7.30 (d, 2H, J = 8.5 Hz), 6.90 (d, 2H, J = 8.5 Hz), 6.58 (d, 1H, J = 5.8 Hz), 3.8 (s, 3H), 2.3 (s, 6H).


Anal Calc'd for C₃₁H₃₄N₇O₂F₃: C 62.72, H 5.77, N 16.52
Found: C 62.57, H 5.95, N 16.76

20

-64-

EXAMPLE 21[4-[2-(1-BENZYLPIPERIDIN-4-YL)-5-(3-TRIFLUOROMETHYLPHENYL)-3H-IMIDAZOL-4-YL]-PYRIMIDIN-2-YL]-4-METHOXYBENZYLAMINE.

5



To a well stirred solution of 4-methoxybenzyl-[4-[2-piperidin-4-yl-5-(3-trifluoromethylphenyl)-3H-imidazol-4-yl]-pyrimidin-2-yl]amine (Example 1H) (100 mg, 0.197 mmole) and benzaldehyde (22 mg, 0.21 mmole) in dichloromethane under argon was added sodium triacetoxyborohydride (44 mg, 0.21 mmole). The mixture was stirred for 4 hours and then quenched by addition of 1N HCl (5 mL). Aqueous sodium bicarbonate (20 mL) and ethyl acetate (20 mL) were added and the layers mixed well. The ethyl acetate layer was removed, concentrated *in vacuo* and residue chromatographed on silica using methylene chloride/methanol/aqueous ammonium hydroxide (97:3:1) to give upon concentration of product containing fractions 94 mg of an oil. Crystallization from ether/hexane (4 mL) gave 80 mg solid.

20

¹H NMR (CDCl₃) δ 9.75 (brs, 1H), 8.10 (d, 1H, J = 5.5 Hz), 7.90 (s, 1H), 7.80 (d, 1H, J = 8.7 Hz), 7.62 (d, 1H, J = 8.7 Hz), 7.54 (t, 1H, J = 8.7 Hz), 7.25-7.4 (m, 7H), 6.90 (d, 2H, J = 8.7 Hz), 6.58 (d, 1H, J = 5.5 Hz), 4.58 (d, 2H, 6.2 Hz), 3.78 (s, 3H), 3.53 (s, 2H), 3.0 (m, 2H), 2.85 (m, 1H), 1.8-2.2 (m, 6H).

25


-65-

Anal Calc'd for C₃₄H₃₃N₆O₃: C 68.21, H 5.56, N 14.04

Found: C 67.99, H 5.59, N 13.88

EXAMPLE 22

5 [4-[2-PIPERIDIN-4-YL-5-(3-TRIFLUOROMETHYLPHENYL)-3H-
IMIDAZOL-4-YL]-PYRIMIDIN-2-YL]AMINE
TRIHYDROCHLORIDE



10


4-Methoxybenzyl-[4-[2-piperidin-4-yl-5-(3-trifluoromethylphenyl)-3H-imidazol-4-yl]pyrimidin-2-yl]amine (Example 1H) (0.125 mg, 0.26 mmole) and 3N HCl (35 mL) were heated to 100°C for 12 hours. The reaction was cooled, washed with diethyl ether (10 mL), and concentrated *in vacuo* to a solid. Trituration from 80% ether/ethanol (10 mL) gave the title compound, 0.102 g, as a yellow solid.

15

MP - 225-230°C

20 ¹H NMR (CD₃OD) δ 8.19 (d, 1H, J = 6.6 Hz), 8.0 (s, 1H), 7.96 (d, 1H, J = 8.4 Hz), 7.89 (d, 1H, J = 8.4 Hz), 7.77 (t, 1H, J = 8.4 Hz), 7.02 (d, 1H, J = 6.6 Hz), 3.2-3.6 (m, 5H), 2.1 = 2.4 (m, 4H).

-66-

EXAMPLE 234-[5-(2-AMINOPYRIMIDIN-4-YL)-1-METHYL-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDINE-1-CARBOXYLIC ACID BENZYL ESTERStep 23A4-[5-(2-Methylthiopyrimidin-4-yl)-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester

To a stirring solution of 4-[5-(2-methylthiopyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1E) (4.0 g, 7.22 mmole) in toluene (80 mL) was added dimethylformamide dimethyl acetal (4.0 mL) and the mixture heated to reflux for 18 hours. The reaction was cooled, concentrated *in vacuo* to a foam and chromatographed on silica using 5% acetone/methylene chloride to give upon concentration of the product containing fractions the title compound 2.78 g.

¹H NMR (CDCl₃) δ 8.33 (d, 1H, J = 6.0 Hz), 7.78 (s, 1H), 7.5-7.6 (m, 2H), 7.3-7.45 (m, 6H); 6.76 (d, 1H, J = 6.0 Hz), 5.13 (s, 2H), 4.35 (brs, 2H), 3.80 (s, 3H), 2.98 (m, 3H), 2.60 (s, 3H), 2.0 (m, 4H).

Additional elution of the above column chromatography with 5% methanol/methylene chloride gave 4-[4-(2-methylthiopyrimidin-4-yl)-1-methyl-5-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester.

-67-

¹H NMR (CDCl₃) δ 8.39 (d, 1H, J = 6.0 Hz), 7.5-7.74 (m, 5H), 7.3 - 7.44 (m, 5H), 5.18 (s, 2H), 4.35 (brs, 2H), 3.38 (s, 3H), 2.7-3.1 (m, 3H), 1.9-2.1 (m, 4H), 1.7 (s, 3H).

5

Step 23B

4-[5-(2-Methylsulfonylpyrimidin-4-yl)-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester

10

The title compound was prepared by oxidation of 4-[5-(2-methylthiopyrimidin-4-yl)-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester as described in Example 1 step F.

15

¹H NMR (CDCl₃) δ 8.58 (d, 1H, J = 5.7 Hz), 7.76 (s, 1H), 7.60 (m, 2H, J = 10.0 Hz), 7.48 (t, 1H, J = 10.0 Hz), 7.37 (m, 5H), 7.24 (d, 1H, J = 5.7 Hz), 5.14 (s, 2H), 4.34 (brs, 2H), 3.98 (s, 3H), 3.40 (s, 3H), 3.01 (m, 3H), 1.90 - 2.0 (m, 4H).

20

Step 23C

4-[5-(2-Aminopyrimidin-4-yl)-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester

25

The title compound was prepared from 4-[5-(2-Methylsulfonylpyrimidin-4-yl)-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester as described in Example 12.

30

MP - 80-83°C

Mass Spectral Analysis - M⁺1 = 537

¹H NMR (CDCl₃) δ 8.18 (d, 1H, J = 5.9 Hz), 7.82 (s, 1H), 7.61 (d, 1H, J = 8.6 Hz), 7.49 (d, 1H, J = 8.6 Hz), 7.3-7.4 (m, 6H), 6.48 (d, 1H, J = 5.9 Hz), 5.1-5.3 (m, 4H), 4.35 (brs, 2H), 3.74 (s, 3H), 2.95 (m, 3H), 1.8-2.1 (m, 4H).


35

-68-

EXAMPLE 24

4-[5-(2-AMINOPYRIMIDIN-4-YL)-1-METHYL-4-(3-
TRIFLUOROMETHYLPHENYL)
-1H-IMIDAZOL-2-YL]-PIPERIDINE

5



The title compound was prepared from 4-[4-(2-Aminopyrimidin-4-yl)-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester as described in
10 Example 1 step H, and isolated as the trifluoroacetic acid salt after purification by preparative HPLC and lyophilization of the product containing fractions.


15 MP = 57-60°C
 ^1H NMR (CD_3OD) 8.06 (d, 1H, $J=6.5$ Hz), 7.84 (s, 1H), 7.72 (m, 2H),
7.62 (t, 1H, $J=8.1$ Hz), 6.53 (d, 1H, $J=6.5$ Hz), 4.00 (s, 3H), 3.2-3.4 (m,
5H), 2.0-2.2 (m, 4H)
20 Mass Spectral Analysis - $\text{M}^+1 = 403$

-69-

EXAMPLE 25

4-[4-(2-AMINOPYRIMIDIN-4-YL)-1-METHYL-5-(3-
TRIFLUOROMETHYLPHENYL)
-3H-IMIDAZOL-2-YL]PIPERIDINE-1-CARBOXYLIC ACID BENZYL
ESTER

5



10 The title compound was prepared from 4-[4-(2-methylthiopyrimidin-4-yl)-1-methyl-5-(3-trifluoromethylphenyl)-3H-imidazol-2-yl]piperidine-1-carboxylic acid benzyl ester as described in Example 12.

MP 215-217°C

15 Anal Calc'd for C₂₈H₂₇N₆O₂F₃: C 62.68, H 5.07, N 15.66
Found: C 62.34, H 4.77, N 15.50


Mass Spectral Analysis - M⁺ = 537

-70-

EXAMPLE 26

4-[4-(2-AMINOPYRIMIDIN-4-YL)-1-METHYL-5-(3-
TRIFLUOROMETHYLPHENYL)
-3H-IMIDAZOL-2-YL]-PIPERIDINE

5




10 The title compound was prepared from 4-[4-(2-aminopyrimidin-4-yl)-1-methyl-5-(3-trifluoromethylphenyl)-3H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester as described in Example 1 step H.

Mass Spectral Analysis - $M^+1 = 403$.

-71-

EXAMPLE 274-([2-PIPERIDIN-4-YL]-5-(3-TRIFLUOROMETHYLPHENYL)-3H-
IMIDAZOL-4-YL)-PYRIMIDINE-2-CARBOXAMIDE
TRIHYDROBROMIDE

5

Step 27A

10 4-[5-[2-Cyanopyrimidin-4-yl]-4-(3-trifluoromethyl)-1H-imidazol-2-yl]-
piperidine-1-carboxylic acid benzyl ester.

15 To a stirred solution of 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 1F) (0.50 g, 0.85 mmole) in dimethyl sulfoxide (3 mL) was added sodium cyanide (0.087 g, 1.70 mmole) and the resulting mixture warmed to 60°C for 2 hours. The reaction was cooled, diluted with water (30 mL) and extracted with ethyl acetate (2x30mL). The ethyl acetate extracts were dried over anhydrous sodium sulfate, filtered and concentrated to an oil. The residue was
20 chromatographed on silica using ethyl acetate/hexane to give upon concentration of the product containing fractions a foam, (0.455 g).

Mass Spectral Analysis - $M^{+1} = 533$.

-72-


Step 27B4-[2-Piperidin-4-yl]-5-(3-trifluoromethylphenyl)-3H-imidazol-4-yl)-pyrimidine-2-carboxamide trihydrobromide

5 The title compound was prepared from 4-[5-[2-cyanopyrimidin-4-yl]-4-(3-trifluoromethyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester as described in Example 1 step H.

10 Mass Spectral Analysis - $M^+1 = 417$
MP - 235-240°C

EXAMPLE 28


15 4-[(2-PIPERIDIN-4-YL)-5-(3-TRIFLUOROMETHYLPHENYL)-3H-IMIDAZOL-4-YL]-PYRIMIDINE



20 4-[5-[2-Cyanopyrimidin-4-yl]-4-(3-trifluoromethyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (0.05 g, 0.09 mmole) and 3N HCl (3 mL) were heated to 100°C for 72 hours. The reaction was cooled and concentrated to a solid. The solid was purified by preparative HPLC to give the title compound (40 mg) upon lyophilization of the product containing fractions.

25 Mass Spectral Analysis - $M^+1 = 374$.

-73-

EXAMPLE 294-[[2-(PIPERIDIN-4-YL)-5-(3-TRIFLUOROMETHYLPHENYL)-3H-
IMIDAZOL-4-YL]]-PYRIMIDIN-2-CARBOXYLIC ACID


To a stirred solution of 4-[[2-piperidin-4-yl]-5-(3-trifluoromethylphenyl)-3H-imidazol-4-yl]pyrimidine-2-carboxamide trihydrobromide (0.120 g, 0.18 mmole) in methanol (1 mL) was added
10 5N sodium hydroxide (1 mL) and the reaction stirred for 18 hours. The reaction was diluted with 5N HCl (1 mL) and purified by preparative HPLC to give the title compound (100 mg) upon lyophilization of the product containing fractions.

15 Mass Spectral Analysis - $M^{+1} = 418$
MP 60-67°C

-74-

EXAMPLE 301-[4-[5-(2-AMINOPYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDIN-4-YL]ETHANONE

5

Step 30A4-[5-(2-Methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]piperidine

10

To a stirred solution of 4-[5-[2-methylsulfonylpyrimidin-4-yl]-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]piperidine-1-carboxylic acid benzyl ester (Example 1F) (400 mg, 0.683 mmole) in methylene chloride (6 mL), under argon was added dropwise 30% hydrogen bromide /acetic acid (6.0 mL). The reaction was allowed to stir for 1 hour, diluted with diethyl ether (60 mL) and the resulting solid filtered to give 0.474 g of the dihydrobromide salt.

20

MP 125-130°C

Mass Spectral Analysis - $M+1 = 452$.Step 30B1-[4-[5-(2-Methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidin-4-yl]ethanone

25

The title compound was prepared from 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]piperidine as described in Example 19.

-75-

¹H NMR (CD₃OD) Rotamer d 8.6-8.84 (brn, 1H), 7.6-8.2 (m, 5H), 4.60 (brd, 1H, J = 18 Hz), 4.10 (d, 1H, J = 18 Hz), 2.7-3.4 (m, 6H), 2.26 (s, 3H), 1.8-2.2 (m, 4H).

5

Step 30C

1-[4-[5-(2-Aminopyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]piperidin-4-yl]ethanone

10

The title compound was prepared from 1-[4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]piperidin-4-yl]ethanone described in Example 12.

15


¹H NMR (CD₃OD) d 8.14 (d, 1H, J = 6.6 Hz), 7.94 (s, 1H), 7.91 (d, 1H, J = 8.5 Hz), 7.83 (d, 1H, J = 8.5 Hz), 7.73 (t, 1H, J = 8.5 Hz), 6.94 (d, 1H, J = 6.6 Hz), 4.65 (d, 1H, J = 18 Hz), 4.10 (d, 1H, J = 18 Hz), 2.80 (m, 1H), 1.8-2.2 (m, 7H).

20

Anal Calc'd for C₂₁H₂₁N₆OF₃•2.0 CF₃COOH•1.0 H₂O: C 44.37, H 3.72, N 12.42
Found: C 44.16, H 3.51, N 12.53

EXAMPLE 31

25 1-[4-[5-(2-AMINOPYRIMIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-2-YL]-PIPERIDIN-1-YL]PHENYLMETHANONE



-76-

The title compound was prepared from 4-[5-(2-methylsulfonylpyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]piperidine (Example 30A) as described in Examples 19 (replacing acetic acid with benzoic acid) and 12


¹H NMR (CD₃OD) δ 8.14 (d, 1H, J = 6.5 Hz), 7.95 (s, 1H), 7.92 (d, 1H, J = 8.7 Hz), 7.84 (d, 1H, J = 8.7 Hz), 7.73 (t, 1H, J = 8.7 Hz), 7.4-7.6 (m, 5H), 6.96 (d, 1H, J = 6.5 Hz), 3.85 (brm, 1H), 3.05 (m, 1H), 1.8-2.2 (m, 6H).

Anal Calc'd for C₂₆H₂₃N₆OF₃•2CF₃COOH, 0.5H₂O: C 49.28, H 3.54, N 11.84

Found: C 49.38, H 3.59, N 11.52

EXAMPLE 32

N-([5-(2-PIPERIDIN-4-YL)-4-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-4-YL]PYRIMIDIN-2-YL)}ACETAMIDE



Step 32A

4-[5-(2-Acetamidopyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]piperidine-1-carboxylic acid benzyl ester

-77-

To a stirred solution of 4-[5-(2-aminopyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (Example 12) (0.220 g, 0.421 mmole) in THF (4 mL) under argon at 0°C was added diisopropylethylamine (0.22 mL, 1.26 mmole) followed by acetyl chloride (0.036 mL, 0.50 mmole). The reaction was stirred at 20°C for 1 hour, diluted with water (10 mL) and product extracted with ethyl acetate (2x50 mL). The ethyl acetate extracts were dried over anhydrous sodium sulfate, filtered, concentrated and chromatographed on silica using 5% methanol/methylene chloride to give the product (0.170 g) as an oil.

¹H NMR (CDCl₃) Rotamers d 10.2 (brs, 1H), 8.56 (d, 1H, J = 6.6 Hz), 7.2-8.0 (m, 10H), 5.12 (brs, 2H), 4.30 (brs, 2H), 2.9-3.1 (m, 3H), 2.32 (s, 3H), 1.8-2.2 (m, 5H).

Example 32B

N-[5-(2-Piperidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-4-yl]pyrimidin-2-yl]acetamide


The title compound was prepared from 4-[5-(2-acetamidopyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]piperidine-1-carboxylic acid benzyl ester as described in Example 30, Step A.

Mass Spectral Analysis M⁺1 = 431
MP >200°C (Dec)

-78-

EXAMPLE 33

N-[4-(2-PIPERIDIN-4-YL)-5-(3-TRIFLUOROMETHYLPHENYL)-1H-IMIDAZOL-4-YL]-PYRIMIDIN-2-YL]BENZAMIDE



5

The title compound was prepared from 4-[5-(2-aminopyrimidin-4-yl)-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester as described in Example 32 (replacing acetyl chloride with) benzoyl chloride.

10

MP >260°C (Dec)


Mass Spectral Analysis - $M^+1 = 493$

-79-

EXAMPLE 34

(S)-4-[5-(3-METHYL-2-PIPERIDIN-4-YL-5-(3-TRIFLUOROMETHYLPHENYL)-3H-IMIDAZOL-4-YL)-PYRIMIDIN-2-YL]-1-PHENYLETHYL)AMINE

5

STEP 34A

10 4-[5-(2-Methylsulfonylpyrimidin-4-yl)-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester

The title compound was prepared using the procedure set forth in Example 1, Steps 1A to 1E, and Example 23, Steps 23A and 23B.

15

STEP 34B

20 (S)-4-[5-(2-(1-Phenylethylamino)pyrimidin-4-yl)-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester

25 A mixture of 4-[5-(2-methylsulfonylpyrimidin-4-yl)-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (12.6 gm, 0.021 mole) and S(-)-(a)-methylbenzylamine (25.0 gm, 0.21 mole) was heated to 100°C for 1 hour, under argon. The mixture was cooled and chromatographed on silica (1 kg) using 40% ethyl acetate/hexane to give (S)-4-[5-[2-(1-

-80-

phenylethylamino)-pyrimidin-4-yl]-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (13.4 gms.)

5

STEP 34C

(S)-4-[5-(2-(1-Phenylethylamino)-pyrimidin-4-yl)-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]piperidine

To a solution of (S)-4-[5-[2-(1-phenylethylamino)-pyrimidin-4-yl]-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (12.5 gm, 0.0195 mole) in dichloromethane (170 mL) at 0 °C, under argon, was slowly added 30 % hydrogen bromide in acetic acid (170 mL). The solution was allowed to stir for 1.5 h at 0°C and then diluted with diethyl ether (2.0 L). The resulting solid was filtered under argon, washed with diethyl ether (500 ml) and sucked dry under argon. The solid was then dissolved in dichloromethane (500 mL) and 10 % aqueous sodium bicarbonate (500 mL) and mixed well. The dichloromethane solution was removed and the aqueous layer washed with dichloromethane (200 mL). The dichloromethane extracts were dried over anhydrous sodium sulfate, concentrated and the resulting foam (9.6 gm) chromatographed on silica using (dichloromethane/methanol/acetic acid/water- 90/10/1/1). The resulting product containing fractions were washed with 10 % aqueous sodium bicarbonate, dried over anhydrous sodium sulfate and concentrated to give (S)-4-[5-[2-(1-phenylethylamino)-pyrimidin-4-yl]-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine (9.4 gm). The solid was dissolved in ethyl acetate (400 ml) filtered and then treated with a solution of hydrochloric acid/ethyl acetate (84 mL, 0.033 gm HCl/mL, 4 eq.). The resulting solid was filtered, dissolved in water (100 ml) and lyophilized to give (S)-4-[5-[2-(1-phenylethylamino)-pyrimidin-4-yl]-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine trihydrochloride (12.0 gm).

Anal. Calc. for C₂₈H₂₉N₆F₃ .3.0 HCl 2.7 H₂O. C, 50.57 ; H, 5.60 ; N, 12.57


-81-

Found C, 50.39; H, 5.47; N, 12.33

¹H NMR (CD₃OD) δ 8.27 (brs, 1H), 8.0 (s, 1H), 7.91 (d, 1H, J = 8.7 Hz), 7.82 (d, 1H, J = 8.7 Hz), 7.73 (t, 1H, J = 8.7 Hz), 7.2-7.6 (m, 5H), 6.64 (d, 1H, J = 6.5 Hz), 5.2 (brs, 1H), 3.2-4.0 (m, 8H), 2.2-2.4 (m, 4H), 1.65 (d, 3H, J = 7.2 Hz).

EXAMPLE 35

10 (S)-[4-[3-METHYL-2-(1-METHYLPYPERIDIN-4-YL)-5-(3-TRIFLUOROMETHYLPHENYL)-3H-IMIDAZOL-4-YL]-PYRIMIDINE-2-YL]-(1-PHENYLETHYL)AMINE




15 The title compound was prepared from (S)-4-[5-(2-(1-phenylethylamino)pyrimidin-4-yl)-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester as described in example 18.

20 Anal. Calc'd for C₂₉H₃₁N₆F₃ 3CF₃COOH: C 48.73, H 3.97, N 9.74
Found: C 48.92, H 4.20, N 9.85

-82-

EXAMPLE 36CYCLOPROPYL-[4-[3-METHYL-2-(PIPERIDIN-4-YL)-5-(3-TRIFLUOROMETHYLPHENYL)-3H-IMIDAZOL-4-YL]-PYRIMIDIN-2-YL]AMINE

5



10 The title compound was prepared from 4-[5-(2-methylsulfonylpyrimidin-4-yl)-1-methyl-4-(3-trifluoromethylphenyl)-1H-imidazol-2-yl]-piperidine-1-carboxylic acid benzyl ester (example 23, step B) as described in example 34, steps B (replacing S(-)-(a)-methybenzylamine with cyclopropylamine) and C.


15 Anal Calc'd C₂₃H₂₅N₆F₃ 3.0 CF₃COOH 1.0 H₂O: C 43.40, H 3.77, N 10.47
Found: C 43.41, H 3.74, N 10.53

-83-

EXAMPLE 37

(S)-[4-[3-METHYL-2-(1-CYCLOPROPYLPYPERIDIN-4-YL)-5-(3-TRIFLUOROMETHYL
PHENYL)-3H-IMIDAZOL-4-YL]-PYRIMIDINE-2-YL]-(1-PHENYLETHYL)AMINE

5



To a solution of (S)-[4-[3-methyl-2-piperidin-4-yl-5-(3-trifluoromethylphenyl)-3H-imidazol-4-yl]-pyrimidin-2-yl]-(1-phenylethyl)amine (300 mg, 0.592 mmole) in methanol (5 ml) under argon was added acetic acid (0.34 ml, 6.0 mmole) and [(1-ethoxycyclopropyl)oxy]-trimethylsilane (0.6 ml, 3.0 mmole). After stirring for 5 minutes sodium cyanoborohydride (150 mg, 2.30 mmole) was added. The mixture was warmed to reflux for 4 hours cooled and concentrated to an oil. The oil was dissolved in methylene chloride (30 ml) and water (20 ml) added. The pH was adjusted to 10.0 with 1N sodium hydroxide and the methylene chloride layer removed, dried over sodium sulfate and concentrated to a foam. The foam was purified by reverse phase preparative HPLC chromatography using 0.1% trifluoroacetic acid/water and acetonitrile to give upon lyophilization of the product containing fractions 480 mg of the title compound.


Anal Calc'd for $C_{31}H_{33}N_6F_3$ 3.0 CF_3COOH : C 50.00, H 4.08, N 9.46
 Found: C 50.39, H 4.24, N 9.76

-84-

EXAMPLE 38

(S)-[4-[2-(1-CYCLOPROPYLMETHYL-PIPERIDIN-4-YL)-3-
METHYL-5-(3-TRIFLUOROMETHYLPHENYL)-3H-IMIDAZOL-4-
YL]-PYRIMIDIN-2-YL]-(1-PHENYLETHYL)-AMINE
TRIHYDROCHLORIDE

5



10 The title compound was prepared from (S)-[4-[3-methyl-2-piperidin-4-yl-5-(3-trifluoromethylphenyl)-3H-imidazol-4-yl]-pyrimidin-2-yl]-(1-phenylethyl)amine (example 34) as described in example 21 (replacing benzaldehyde with cyclopropanecarboxaldehyde).


15 Anal Calc'd for C₃₂H₃₅N₆F₃ 3.0HCl 2.55 H₂O: C 53.68, H 6.07, N 11.74
Found: C 53.79, H 6.24, N 11.34
M.P. 145-155°C

-85-

EXAMPLE 39

(S)-(1-PHENYLETHYL)-{4-[2-PIPERIDIN-4-YL-3-PROPYL-5-(3-TRIFLUOROMETHYLPHENYL)-3H-IMIDAZOL-4-YL]-PYRIMIDIN-2-YL}AMINE TRIHYDROCHLORIDE

5



10 The title compound was prepared using the procedures set forth in Example 1, Steps 1A to 1E, Example 23, Steps 23A (replacing dimethylformamide dimethyl acetal with dimethylformamide dipropyl acetal) and 23B and Example 34, Steps 34B and 34C.


Anal Calc'd for $C_{30}H_{33}N_6F_3 \cdot 3.0HCl \cdot 0.7 H_2O$: C 54.87, H 5.74, N 12.80

15 Found: C 54.92, H 5.98, N 12.55

-86-

EXAMPLE 40

{4-[5-(3,4-DICHLOROPHENYL)-3-METHYL-2-PIPERIDIN-4-YL-3H-IMIDAZOL-4-YL]-PYRIMIDIN-2-YL}-(1-PHENYLETHYL)-AMINE



5

The title compound was prepared using the procedures set forth in Example 1, Steps 1A to 1E (replacing N-methoxy-N-methyl-3-trifluoromethylbenzamide with N-methoxy-N-methyl-3,4-dichlorobenzamide), Example 23, Steps 23A and 23B and Example 34, Steps 34B and 34C.

10

Anal Calc'd for $C_{27}H_{28}N_6Cl_2 \cdot 0.6 CH_3Cl_2$: C 59.36, H 5.27, N 15.05
Found: C 15.52, H 5.07, N 15.03


15

-87-

EXAMPLE 41

[4-[5-(3,4-DICHLOROPHENYL)-3-METHYL-2-PIPERIDIN-4-YL-3H-
IMIDAZOL-4-YL]-PYRIMIDIN-2-YL]-(4-METHOXYBENZYL)-
AMINE

5



10 The title compound was prepared using the procedures set forth in Example 1, Steps 1A to 1E (replacing N-methoxy-N-methyl-3-trifluoromethylbenzamide with N-methoxy-N-methyl-3,4-dichlorobenzamide), Example 23, Steps 23A and 23B and Example 34, Steps 34B (replacing S-(a)-methylbenzylamine with 4-methoxybenzylamine) and 34C.


15 Anal Calc'd for C₂₇H₂₈N₆Cl₂O • 0.60 CH₂Cl₂: C 57.71, H 5.12, N 14.63

Found: C 57.90, H 5.24, N 14.36

-88-

EXAMPLE 42[4-[5-(3,4-DICHLOROPHENYL)-3-PROPYL-2-PIPERIDIN-4-YL-3H-IMIDAZOLE-4-YL]PYRIMIDIN-2-YL]-(1-PHENYLETHYL)-AMINE

5




The title compound was prepared using the procedures set forth in Example 1, Steps 1A to 1E (replacing N-methoxy-N-methyl-3-trifluoromethylbenzamide with N-methoxy-N-methyl-3,4-dichlorobenzamide), Example 23, Steps 23A (replacing dimethylformamide dimethyl acetal with dimethylformamide dipropyl acetal) and 23B and Example 34, Steps 34B and 34C.

¹H NMR (CDCl₃): d (8.14, J=5.13Hz, d, 1H); (7.64, s, 1H); (7.19 - 7.40, m, 8H); (6.37, J=4.88Hz, d, 1H); (5.78, bs, 1H); (5.21, J=6.84Hz, 13.92, q, 1H); 3.91 - 4.08, bm, 1H); (3.60 - 3.78, m, 3H); (3.47, J=6.6Hz, q, 1H); (3.30 - 3.36, m, 2H); (2.78 - 2.86, m, 3H); (1.86 - 1.98, m, 2H); (1.58, J=6.84,Hz d, 3H); (1.00 - 1.53, m, 3H); (0.60, bs,2H).

20

-89-

EXAMPLE 43CYCLOPROPYL-[4-[5-(3,4-DICHLOROPHENYL)-2-PIPERIDIN-4-
YL-3-PROPYL-3H-IMIDAZOL-4-YL]-PYRIMIDIN-2-YL]-AMINE


5

The title compound was prepared using the procedures set forth in Example 1, Steps 1A to 1E (replacing N-methoxy-N-methyl-3-trifluoromethylbenzamide with N-methoxy-N-methyl-3,4-dichlorobenzamide), Example 23, Steps 23A (replacing dimethylformamide dimethyl acetal with dimethylformamide dipropyl acetal) and 23B and Example 34, Steps 34B (replacing S-(a)-methylbenzylamine with cyclopropylamine) and 34C.

15 ¹H NMR (CDCl₃): d (8.19, J=4.64Hz, bd, 1H); (7.68, s, 1H); (7.24 - 7.33, b, 2H); (6.44, J=5.12Hz, bd, 1H); (5.47, s, 1H); (4.26, bs, 2H); (3.30, J=12.69Hz, bd, 2H); (2.76 - 2.88, m, 4H); (2.56, bs, 2H); (1.88 - 2.17, m, 4H); (1.57 - 1.64, m, 2H); (1.25, s, 1H); (0.83 - 0.98, m, 2H); (0.60 - 0.66, bs, 2H).

20

-90-

EXAMPLE 44CYCLOHEXYL-[4-[5-(3,4-DICHLOROPHENYL)-2-PIPERIDIN-4-YL-3-PROPYL-3H-IMIDAZOL-4-YL]-PYRIMIDIN-2-YL]-AMINE

5

- The title compound was prepared using the procedures set forth in Example 1, Steps 1A to 1E (replacing N-methoxy-N-methyl-3-trifluoromethylbenzamide with N-methoxy-N-methyl-3,4-dichlorobenzamide), Example 23, Steps 23A (replacing dimethylformamide dimethyl acetal with dimethylformamide dipropyl acetal) and 23B and Example 34, Steps 34B (replacing S-(a)-methylbenzylamine with cyclohexylamine) and 34C.
- ¹H NMR (CDCl₃): d (8.14, J=4.88Hz, bd, 1H); (7.63, s, 1H); (7.23 - 7.32, m, 2H); (6.35, J=6.1Hz, d, 1H); (5.13, J=7.81Hz, bd, 1H); (4.14 - 4.19, m, 2H); (3.81 - 3.91, m, 1H); (3.32, J=12.7Hz, bd, 2H); (2.71 - 2.97, m, 4H); (2.17, s, 2H); (1.01 - 2.17, m, 15H); (0.90 - 0.99, m, 3H).

- The ability of compounds of the present invention to inhibit the synthesis or the activity of cytokines can be demonstrated using the following *in vitro* assays.

Lipopolysaccharide mediated production of cytokines

- Human peripheral blood mononuclear cells (PBMC) are isolated from fresh human blood according to the procedure of Chin and

25

-91-

Kostura, *J. Immunol.* **151**, 5574-5585 (1993). Whole blood is collected by sterile venipuncture into 60 mL syringes coated with 1.0 mL of sodium- heparin (Upjohn, 1000 U/mL) and diluted 1:1 in Hanks Balanced Salt Solution (Gibco). The erythrocytes are separated from the PBMC's by centrifugation on a Ficoll-Hypaque lymphocyte separation media. The PBMC's are washed three times in Hanks Balanced Salt Solution and then resuspended to a final concentration of 2×10^6 cell/mL in RPMI containing 10% fresh autologous human serum, penicillin streptomycin (10 U/mL) and 0.05% DMSO. Lipopolysaccharide (Salmonella type Re545; Sigma Chemicals) is added to the cells to a final concentration of 100 ng/mL. An aliquot (0.1 mL) of the cells is quickly dispensed into each well of a 96 well plate containing 0.1 mL of the test compound, at the appropriate dilution, and are incubated for 24 hours. at 37°C in 5% CO₂. At the end of the culture period, cell culture supernatants are assayed for IL-1b, TNF-a, IL-6 and PGE₂ production using specific ELISA.

IL-1 mediated cytokine production

Human peripheral blood mononuclear cells are isolated from fresh human blood according to the procedure of Chin and Kostura, *J. Immunol.* **151**, 5574-5585 (1993). Whole blood is collected by sterile venipuncture into 60 mL syringes coated with 1.0 mL of sodium- heparin (Upjohn, 1000 U/mL) and diluted 1:1 in Hanks Balanced Salt Solution (Gibco). The erythrocytes are separated from the PBMC's by centrifugation on a Ficoll-Hypaque lymphocyte separation media. The PBMC's are washed three times in Hanks Balanced Salt Solution and then resuspended to a final concentration of 2×10^6 cell/mL in RPMI containing 10% fresh autologous human serum, penicillin streptomycin (10 U/mL) and 0.05% DMSO. Endotoxin free recombinant human IL-1b is then added to a final concentration of 50 pMolar. An aliquot (0.1 mL) of the cells is quickly dispensed into each well of a 96 well plate containing 0.1 mL of the compound at the appropriate dilution. and are incubated for 24 hours. at 37°C in 5% CO₂. At the end of the culture

-92-

period, cell culture supernatants are assayed for TNF-a, IL-6 and PGE₂ synthesis using specific ELISA..

5 Determination of IL-1b , TNF-a, IL-6 and prostanoid production from LPS or IL-1 stimulated PBMC's

IL-1b ELISA

 Human IL-1b can be detected in cell-culture supernatants or whole blood with the following specific trapping ELISA. Ninety-six well
10 plastic plates (Immulon 4; Dynatech) are coated for 12 hours at 4°C with 1 mg/mL protein-A affinity chromatography purified mouse anti-human IL-1b monoclonal antibody (purchased as an ascites preparation from LAO Enterprise, Gaithersburg Maryland.) diluted in Dulbecco's phosphate-buffered saline (-MgCl₂, -CaCl₂). The plates are washed with
15 PBS-Tween (Kirkegaard and Perry) then blocked with 1% BSA diluent and blocking solution (Kirkegaard and Perry) for 60 minutes at room temperature followed by washing with PBS Tween. IL-1b standards are prepared from purified recombinant IL-1b produced from *E. coli*.. The highest concentration begins at 10 ng/mL followed by 11 two-fold serial
20 dilutions. For detection of IL-1b from cell culture supernatants or blood plasma, 10 - 25 mL of supernatant is added to each test well with 75 - 90 mL of PBS Tween. Samples are incubated at room temperature for 2 hours then washed 6 times with PBS Tween on an automated plate washer (Dennly). Rabbit anti-human IL-1b polyclonal antisera diluted
25 1:500 in PBS-Tween is added to the plate and incubated for 1 hour at room temperature followed by six washes with PBS-Tween. Detection of bound rabbit anti-IL-1b IgG is accomplished with Fab' fragments of Goat anti-rabbit IgG-horseradish peroxidase conjugate (Accurate Scientific) diluted 1:10,000 in PBS-Tween. Peroxidase activity was
30 determined using TMB peroxidase substrate kit (Kirkegaard and Perry) with quantitation of color intensity on a 96-well plate Molecular Devices spectrophotometer set to determine absorbance at 450 nM. Samples are evaluated using a standard curve of absorbance versus concentration.

-93-

Four-parameter logistics analysis generally is used to fit data and obtain concentrations of unknown compounds.

TNF-a ELISA

5 Immulon 4 (Dynatech) 96-well plastic plates are coated with
a 0.5 mg/mL solution of mouse anti-human TNF-a monoclonal antibody.
The secondary antibody is a 1:2500 dilution of a rabbit anti-human TNF-
a polyclonal serum purchased from Genzyme. All other operations are
10 identical to those described above for IL-1b. The standards are prepared
in PBS-Tween + 10% FBS or HS. Eleven 2 fold dilutions are made
beginning at 20 ng/mL TNF-a.

IL-6 ELISA

15 Levels of secreted human IL-6 are also determined by
specific trapping ELISA as described previously in Chin and Kostura, *J.*
Immunol. **151**, 5574-5585 (1993). (Dynatech) ELISA plates are coated
with mouse anti-human IL-6 monoclonal antibody diluted to 0.5 mg/mL
in PBS. The secondary antibody, a rabbit anti-human IL-6 polyclonal
20 antiserum, is diluted 1:5000 with PBS-Tween. All other operations are
identical to those described above for IL-1b. The standards are prepared
in PBS-Tween + 10% FBS or HS. Eleven 2 fold dilutions are made
beginning at 50 ng/mL IL-6.

PGE₂ production

25 Prostaglandin E₂ is detected in cell culture supernatants
from LPS or IL-1 stimulated PBMC's using a commercially available
enzyme immunoassay . The assay purchased from the Cayman Chemical
(Catalogue number 514010) and is run exactly according to the
manufacturers instructions.

30

Interleukin8 (IL-8)

The present compounds can also be assayed for IL-8
inhibitory activity as discussed below. Primary human umbilical cord
endothelial cells (HUVEC) (Cell Systems, Kirland, Wa) are maintained


-94-

5 in culture medium supplemented with 15% fetal bovine serum and 1 %
CS-HBGF consisting of aFGF and heparin. The cells are then diluted 20-
fold before being plated (250 μ l) into gelatin coated 96-well plates. Prior
to use, culture medium is replaced with fresh medium (200 μ l). Buffer or
10 test compound (25 μ l, at appropriate concentrations) is then added to each
well in quadruplicate wells and the plates incubated for 6h in a
humidified incubator at 37°C in an atmosphere of 5% CO₂. At the end of
the incubation period, supernatant is removed and assayed for IL-8
concentration using an IL-8 ELISA kit obtained from R&D Systems
10 (Minneapolis, MN). All data is presented as mean value (ng/mL) of
multiple samples based on the standard curve. IC₅₀ values where
appropriate are generated by non-linear regression analysis.

-95-

WHAT IS CLAIMED IS:

1. A compound represented by formula I:




5

I

or a pharmaceutically acceptable salt thereof, wherein:

- X and X' each independently represent $-(CH_2)_m-Y-(CH_2)_n-$,
 10 wherein m and n represent integers within the range of from 0 - 4, such that the sum of m and n is from 0 - 6; Y represents a member selected from the group consisting of: a direct bond; O; $S(O)_y$, with y equal to 0, 1 or 2; $NR^{q'}$, with $R^{q'}$ as defined below; $C(O)$; $OC(O)$; $C(O)O$; $SO_xNR^{q'}$ with x equal to 1 or 2 and $R^{q'}$ as defined below; $NR^{q'}SO_x$;
 15 $C(O)NR^{q'}$ and $NR^{q'}C(O)$;



represents a 4 to 10 membered non-aromatic heterocycle containing at least one N atom, and optionally containing 1-2 additional N atoms and 0-1 O or S atom;

R^x represents H, C_{1-6} alkyl(R^q)₃, C_{3-8} cycloalkyl, OC_{1-6} alkyl(R^q)₃ or $C(O)C_{1-6}$ alkyl(R^q)₃;

20



each R independently represents a member selected from the group consisting of: halo; hydroxy; C_{1-6} alkyl(R^q)₃; OC_{1-6}

-96-

- alkyl(R^q)₃; C₃₋₈ cycloalkyl(R^q)₃; CN; CONH₂; CONHC₁₋₆ alkyl(R^q)₃; CON(C₁₋₆ alkyl(R^q)₃)₂; NH₂; NHC₁₋₆ alkyl(R^q)₃; NHC₃₋₈ cycloalkyl; N(C₁₋₆ alkyl(R^q)₃)₂; CON(C₃₋₈ cycloalkyl)(C₁₋₆ alkyl(R^q)₃); CO₂H; CO₂C₁₋₆ alkyl(R^q)₃; C(O)C₁₋₆ alkyl(R^q)₃; aryl(R^q)₃; heterocyclyl(R^q)₃; heteroaryl(R^q)₃; CF₃; SH; NO₂; NHSO₂C₁₋₆alkyl(R^q)₃, NHSO₂aryl(R^q)₃, NHSO₂heteroaryl(R^q)₃; N(R^{q'})C(O)C₁₋₆ alkyl(R^q)₃; NR^{q'}C(O)NH(C₁₋₆ alkyl(R^q)₃); C₂₋₄ alkenyl(R^q)₂₋₃ and C₂₋₄ alkynyl(R^q)₁₋₃;
- each R" independently represents a member selected from the group consisting of: halo; hydroxy; C₁₋₆ alkyl(R^q)₃; OC₁₋₆ alkyl(R^q)₃; C₃₋₈ cycloalkyl(R^q)₃; CN; CONH₂; CONHC₁₋₆ alkyl(R^q)₃; CON(C₁₋₆ alkyl(R^q)₃)₂; NH₂; NHC₁₋₆ alkyl(R^q)₃; NHC₃₋₈ cycloalkyl; N(C₁₋₆ alkyl(R^q)₃)₂; CON(C₃₋₈ cycloalkyl)(C₁₋₆ alkyl(R^q)₃); CO₂H; CO₂C₁₋₆ alkyl(R^q)₃; C(O)C₁₋₆ alkyl(R^q)₃; aryl(R^q)₃; heterocyclyl(R^q)₃; heteroaryl(R^q)₃; CF₃; SH; NO₂; SO_yC₁₋₆ alkyl(R^q)₃, with y as defined above; SO₂NH₂; SO₂NHC₁₋₆ alkyl(R^q)₃; SO₂N(C₁₋₆ alkyl(R^q)₃)₂; NHSO₂C₁₋₆alkyl(R^q)₃, NHSO₂aryl(R^q)₃, NHSO₂heteroaryl(R^q)₃; N(R^{q'})C(O)C₁₋₆ alkyl(R^q)₃; NR^{q'}C(O)NH(C₁₋₆ alkyl(R^q)₃); C₂₋₄ alkenyl(R^q)₂₋₃ and C₂₋₄ alkynyl(R^q)₁₋₃;
- each R' independently represents a member selected from the group consisting of: CONH₂; CONHC₁₋₆ alkyl(R^q)₃; CON(C₁₋₆ alkyl(R^q)₃)₂; CONHC₃₋₈ cycloalkyl(R^q)₃; CON(C₃₋₈ cycloalkyl(R^q)₃)₂; CON(C₃₋₈ cycloalkyl)(C₁₋₆ alkyl(R^q)₃); CO₂H; CO₂C₁₋₆ alkyl(R^q)₃; C(O)C₁₋₆ alkyl(R^q)₃; CO₂C₃₋₈ cycloalkyl(R^q)₃; C(O)C₃₋₈ cycloalkyl(R^q)₃; -[C(O)(CH₂)_j-CR⁵R⁶-(CH₂)_k-NR⁷]_p-R⁸; -C(O)C₃₋₈ cycloalkyl(R^q)₃; -C(O)heterocyclyl(R^q)₃; -CON[C₁₋₆alkyl(R^q)₃][C₃₋₈ cycloalkyl(R^q)₃]; -C(O)aryl(R^q)₃; -C(O)heteroaryl(R^q)₃; hydroxy; C₁₋₆ alkyl(R^q)₃; C₃₋₈ cycloalkyl(R^q)₃; OC₁₋₆ alkyl(R^q)₃; OC₃₋₈ cycloalkyl(R^q)₃; heterocyclyl(R^q)₃; CN; NH(R^{q''}); NHC₁₋₆ alkyl(R^q)₃; N(C₁₋₆ alkyl(R^q)₃)₂; NHC₃₋₈ cycloalkyl(R^q)₃; N(C₃₋₈ cycloalkyl(R^q)₃)₂;

-97-

- CF₃; SH; NO₂; C₂₋₄ alkenyl(R^q)₂₋₃, aryl(R^q)₃; heteroaryl(R^q)₃; C₂₋₄ alkynyl(R^q)₁₋₃; -OC(O) C₃₋₈ cycloalkyl(R^q)₃; SO₂NH₂; SO₂NHC₁₋₆ alkyl(R^q)₃; SO₂N(C₁₋₆ alkyl(R^q)₃)₂; NHSO₂C₁₋₆ alkyl(R^q)₃; NHSO₂aryl(R^q)₃; NHSO₂heteroaryl(R^q)₃;
 5 -OC(O)heterocyclyl(R^q)₃; N(R^{q'})C(O)C₁₋₆ alkyl(R^q)₃; NR^{q'}C(O)NH(C₁₋₆ alkyl(R^q)₃); -OC(O)C₁₋₆ alkyl(R^q)₃; -OC(O)aryl(R^q)₃; -OC(O)heteroaryl(R^q)₃; -C(=NR^{q'})NH₂; -C(=NR^{q'})NHC₁₋₆ alkyl(R^q)₃; -C(=NR^{q'})N(C₁₋₆ alkyl(R^q)₃)₂;



- 15 wherein j and k independently represent integers of from 0 - 3;

R⁵ and R⁶ are independently H, aryl, C₁₋₆ alkyl(R^q)₃, or CR⁵R⁶ in combination represents a 3, 4, 5 or 6 membered cycloalkyl or heterocyclyl group, an aryl group or a heteroaryl group;

- 20 p represents 1, 2 or 3, with the proviso that when p represents 1, CR⁵R⁶ represents a 3, 4, 5 or 6 membered cycloalkyl group or a heterocyclyl group, an aryl group or a heteroaryl group, and at least one of j and k is 1, 2 or 3;

- 25 R⁷ and R⁸ are independently H, C₁₋₆ alkyl or aryl;

R⁹ represents H, a negative charge balanced by a positively charged group or a protecting group;

- 30 R^q represents a member selected from the group consisting of: R^{q'}; halo; CN; CO₂H; CO₂C₁₋₄ alkyl; C(O)C₁₋₄ alkyl; NH(R^{q''}); aryl(R^a)₃; heteroaryl(R^a)₃; NHC₁₋₄ alkyl; N(C₁₋₄ alkyl)₂; CONH₂;

-98-

SH ; S(O)_yC₁₋₆ alkyl(R^a)₃; C(O)NHC₁₋₆ alkyl(R^a)₃; C(O)N(C₁₋₆ alkyl(R^a)₃)₂; C₃₋₈ cycloalkyl; NHC(NH)NH₂ ; -heteroalkyl(R^a)₃; -NHC(O)NH₂;



wherein and independently represent mono or bicyclic ring systems, non-aromatic or partially aromatic, containing from 5-10 ring atoms, 1-4 of which are N and 0-1 of which are O or S(O)_y, with y equal to 0, 1 or 2, optionally containing 1-2 carbonyl groups;

each R^a independently represents a member selected from the group consisting of: H, C₁₋₆ alkyl, OC₁₋₆ alkyl, aralkyl, substituted aralkyl, heteroaralkyl, substituted heteroaralkyl, aralkoxy, substituted aralkoxy, halo, hydroxy, CN, CONH₂, CONHC₁₋₆ alkyl, CON(C₁₋₆ alkyl)₂, CO₂H, CO₂C₁₋₆ alkyl, C(O)C₁₋₆ alkyl, phenyl, CF₃, SH, NO₂, SO_yC₁₋₆ alkyl, with y as defined above; SO₂NH₂, SO₂NHC₁₋₆ alkyl, NHSO₂(substituted aryl), NHSO₂(substituted heteroaryl), NHSO₂C₁₋₆alkyl, NHSO₂aryl, NHSO₂heteroaryl, NH₂, NHC₁₋₆ alkyl, N(C₁₋₆ alkyl)₂, NHC(O)C₁₋₆ alkyl, NHC(O)NH(C₁₋₆ alkyl), C₂₋₄ alkenyl and C₂₋₄ alkynyl;

R^{q'} represents H, OH, C₁₋₄ alkyl, -OC₁₋₄ alkyl, aryl or C(O)C₁₋₄ alkyl, and

25

R^{q''} represents H, OH or OC₁₋₄ alkyl.

2. A compound in accordance with claim 1 wherein one or two R'' groups are present, and each independently represents NH₂, NHC₁₋₆ alkyl(R^q)₃, N(C₁₋₆ alkyl)₂, NHC₃₋₈ cycloalkyl,

30

-99-

$N(R^q)C(O)C_{1-6} \text{ alkyl}(R^q)_3$, $C_{1-6} \text{ alkyl}(R^q)_3$, $OC_{1-6} \text{ alkyl}(R^q)_3$, CO_2H , $CONH_2$, $NR^qC(O)NHC_{1-6} \text{ alkyl}(R^q)_3$ or heterocyclyl(R^q)₃.

3. A compound in accordance with claim 1 wherein:
 5 HETCy represents a 5-6 membered non-aromatic heterocycle with 1-2 nitrogen atoms contained therein.

4. A compound in accordance with claim 3 wherein
 10 HETCy represents a pyrrolidinyl or piperidinyl group.

5. A compound in accordance with claim 4 wherein
 HETCy represents a 4-piperidinyl group.


6. A compound in accordance with claim 1 wherein
 15 R' is selected from $C_{1-6} \text{ alkyl}(R^q)_3$, $OC_{1-6} \text{ alkyl}(R^q)_3$, $-C(O)C_{1-6} \text{ alkyl}(R^q)_3$; CN , NO_2 and $CO_2C_{1-6} \text{ alkyl}(R^q)_3$.

7. A compound in accordance with claim 1 wherein
 20 from 1-3 R groups are present and each independently represents a member selected from the group consisting of: halo, hydroxy, $C_{1-6} \text{ alkyl}(R^q)_3$, $OC_{1-6} \text{ alkyl}(R^q)_3$, NH_2 , $NHC_{1-6} \text{ alkyl}(R^q)_3$, $N(C_{1-6} \text{ alkyl}(R^q)_3)_2$ and CF_3 .

8. A compound in accordance with claim 7 wherein
 25 one or two R groups are present, selected from halo and CF_3 .

9. A compound in accordance with claim 1 wherein
 R^x is H , $C_{3-8} \text{ cycloalkyl}$ or $C_{1-6} \text{ alkyl}(R^q)_3$.

30 10. A compound in accordance with claim 9 wherein
 R^x represents H , CH_3 , CH_2CH_3 , $CH_2CH_2CH_3$, $CH_2CH_2CH_2CH_3$,



-100-

11. A compound in accordance with claim 1 wherein X' represents a direct bond.

5 12. A compound in accordance with claim 1 wherein X represents $-(CH_2)_m-Y-(CH_2)_n-$, Y represents a direct bond, O, S or C(O); m represents 0 or 1 and n represents 0 or 1.

10 13. A compound in accordance with claim 12 wherein X represents a direct bond.

14. A compound in accordance with claim 1 wherein one or two R" groups are present, each independently representing NH_2 , $NHC_{1-6} alkyl(R^9)_3$, $N(C_{1-6} alkyl)_2$, $NHC_{3-8} cycloalkyl$, $N(R^9)C(O)C_{1-6} alkyl(R^9)_3$, $C_{1-6} alkyl(R^9)_3$, $OC_{1-6} alkyl(R^9)_3$, CO_2H , $CONH_2$, $NR^9C(O)NHC_{1-6} alkyl(R^9)_3$ or heterocyclyl(R⁹)₃;

HETCy represents a 5-6 membered non-aromatic heterocyclyl with 1-2 nitrogen atoms contained therein;

20 R' is selected from $C_{1-6} alkyl(R^9)_3$, $OC_{1-6} alkyl(R^9)_3$, $-C(O)C_{1-6} alkyl(R^9)_3$, CN; NO₂; and $CO_2C_{1-6} alkyl(R^9)_3$;

25 from 1-3 R groups are present and each independently represents a member selected from the group consisting of: halo, hydroxy, $C_{1-6} alkyl(R^9)_3$, $OC_{1-6} alkyl(R^9)_3$, NH_2 , $NHC_{1-6} alkyl(R^9)_3$, $N(C_{1-6} alkyl(R^9)_3)_2$, CO_2H and CF_3 ;

R^x is H, $C_{3-8} cycloalkyl$ or $C_{1-6} alkyl(R^9)_3$;

X' represents a direct bond;

30 X represents $-(CH_2)_m-Y-(CH_2)_n-$, Y represents a direct bond; m represents 0 or 1 and n represents 0 or 1.

15. A compound in accordance with claim 14 wherein: one or two R" groups are present, selected from the group consisting of: NH_2 , $NHC_{1-6} alkyl(R^9)_3$, $N(C_{1-6} alkyl)_2$, $C_{1-6} alkyl(R^9)_3$, $OC_{1-6} alkyl(R^9)_3$ and heterocyclyl(R⁹)₃;


-101-

HETCy represents a piperidinyl group;
 zero or one R' is present, which is selected from C₁-6
 alkyl(R^q)₃, -CO₂C₁-6 alkyl(R^q)₃ and -C(O)C₁-6 alkyl(R^q)₃;
 one or two R groups are present, selected from halo and

5 CF₃;


R^x is H, C₃-8 cycloalkyl or C₁-6 alkyl(R^q)₃; and
 X and X' represent a direct bond.

16. A compound in accordance with claim 14 wherein:
 one R'' group is present and is selected from:



15 HETCy represents a 4-piperidinyl group;
 R' is absent;
 one or two R groups are present selected from halo and
 CF₃;

R^x represents H, CH₃, CH₂CH₃, CH₂CH₂CH₃,




20




and X and X' represent direct bonds.

-102-

17. A compound represented by one of the structural formulas:




5




10


-103-





5





10









-104-









5









-105-




-106-



-107-




-108-



or a pharmaceutically acceptable salt thereof.

5


18. A compound represented by the formula:



10 or a pharmaceutically acceptable salt thereof.



19. A compound represented by the formula:

-109-







or a pharmaceutically acceptable salt thereof.





- 5 20. A compound represented by one of the structural formulas:



-110-



-111-



5

or a pharmaceutically acceptable salt thereof.

21. A pharmaceutical composition comprised of a compound in accordance with claim 1 in combination with a pharmaceutically acceptable carrier.

10

22. A method of treating a cytokine mediated disease in a mammalian patient in need of such treatment comprising administering to

-112-

said mammal a compound in accordance with claim 1 in an amount which is effective to treat said cytokine mediated disease.

5 23. A method in accordance with claim 22 wherein the cytokine mediated disease is rheumatoid arthritis, osteoarthritis, endotoxemia, toxic shock syndrome, inflammatory bowel disease, tuberculosis, atherosclerosis, muscle degeneration, cachexia, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis or acute synovitis.

10 24. A method in accordance with claim 22 wherein the cytokine mediated disease is rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, adult respiratory distress
15 syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcosis, bone resorption diseases, reperfusion injury, graft v. host rejection, allograft rejection, fever, myalgia due to infection, cachexia secondary to infection or malignancy, cachexia
20 secondary to acquired immune deficiency syndrome (AIDS), AIDS related complex (ARC), keloid formation, scar tissue formation, Crohn's disease, ulcerative colitis or pyresis.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/09888

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :C07D 403/14, 413/14, 417/14

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/256, 269, 272, 274, 275; 544/333, 331, 295, 296, 300, 310, 315, 316, 317, 318, 319, 320, 321, 324, 327, 328

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN, Chemical Abstracts search by structure

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, P	Chemical Abstract. Vol, 126, No. 12, 07 July 1997 (Columbus, OH USA), the abstract No. 126:144274K, ADAMS, J.L. "Imidazole" compounds useful as cytskine inhibitors".	1-24

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" documents which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

29 AUGUST 1997

Date of mailing of the international search report

10 OCT 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JOHN M. FORD

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/09888

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

514/256, 269, 272, 274, 275; 544/333, 331, 295, 296, 300, 310, 315, 316, 317, 318, 319, 320, 321, 324, 327, 328